

Dermatitis artefacta: a challenging diagnosis

Giulia Ciccarese,¹ Giorgia Salvia,² Cristian Fidanzi,^{2,3} Mario Mastrodonato,¹ Francesco Drago⁴

¹Unit of Dermatology, Department of Medical and Surgical Sciences, University of Foggia; ²Unit of Dermatology, Department of Medical and Oncology Area, University of Pisa; ³Unit of Dermatology, Hospital of Carrara; ⁴Section of Dermatology, DISSAL, IRCCS Ospedale Policlinico San Martino, University of Genoa, Italy

Dear Editor,

Dermatitis factitia (DF), also known as dermatitis artefacta, is a rare condition characterized by self-inflicted skin lesions.¹ These lesions can be a result of compulsive habits driven by emotional distress, serving as a means of emotional release during episodes of anxiety.¹ Alternatively, they may be deliberately induced by the patient with the intention of gaining medical attention.¹ DF is essentially the physical manifestation of underlying psychiatric disorders, making it a particularly challenging condition within

Correspondence: Cristian Fidanzi, MD, Unit of Dermatology, University of Pisa, via Roma 67, 56126 Pisa, Italy.
E-mail: cri.fidanzi@outlook.it
Tel.: 340.0630740.

Key words: dermatitis factitia; dermatitis artefacta; psychocutaneous disease; obsessive-compulsive disorder; emotional distress.

Conflict of interest: the authors have no conflict of interest to declare.

Ethics approval and consent to participate: no ethical committee approval was required for this case report by the Department, as it does not contain any studies involving human participants or animals. Informed consent was obtained from the patient included in this study.

Consent for publication: the patient gave her written consent to use her personal data for the publication of this manuscript and any accompanying images.

Availability of data and materials: data supporting the findings of this study are available from the corresponding author upon request.

Received: 7 April 2024.
Accepted: 6 August 2024.

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).

©Copyright: the Author(s), 2025
Licensee PAGEPress, Italy
Dermatology Reports 2025; 17:10014
doi:10.4081/dr.2024.10014

Publisher's note: all claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

the realm of medical practice.² Being a diagnosis primarily based on the exclusion of an underlying disorder often requires multiple investigations that can impose a significant economic burden.³ We present the case of a 70-year-old woman who came to our attention with skin lesions located on her left cheek, left eyebrow, and forehead, which had been present for approximately two years. The patient had a medical history of arterial hypertension, diabetes, and anxious depressive syndrome, for which she was receiving treatment with duloxetine (60 mg daily) and alprazolam (0.25 mg three times daily). On physical examination, the lesions appeared as erythematous and crusted plaques with irregular margins, slightly infiltrated, and were reported as itchy and painful by the patient (Figure 1 a,b). These lesions had proven resistant to the topical antibiotic therapy previously prescribed by another physician. Subsequently, we ordered a comprehensive hematological diagnostic assessment, including blood count, C-reactive protein (CRP), erythrocyte sedimentation rate (ESR), antinuclear antibodies (ANA), extractable nuclear antigen antibodies (anti-ENA), C3, C4, protein electrophoresis, rheumatoid factor, quantiferon, angiotensin-converting enzyme test, thyroid-stimulating hormone, glycemia, glycated hemoglobin, aspartate aminotransferase, alanine aminotransferase, creatinine, and gamma-glutamyl transferase. We started treatment with an emollient. During her second visit to our department, one month after her initial visit, the patient provided her blood test results, which were normal, except for an elevated ESR (25 mm/hr). We recommended a skin biopsy with histological examination and direct immunofluorescence to investigate the possibility of connectivitis, pseudolymphoma, or cutaneous sarcoidosis, which the patient declined. She additionally reported an increase in the dosage of alprazolam (0.50 mg three times a day), prescribed by her psychiatrist for managing her anxious-depressive syndromes. Unexpectedly, after the alprazolam dosage was raised, there was a noticeable improvement in the facial lesions within five months (Figure 1 c,d). However, a persistent erythematous area on the chin remained. Initially, the patient mentioned using tweezers to extract white hair. Upon additional questioning, she revealed that, in the past, she had consistently used tweezers on her face at the level of the previous lesions for hair removal. Notably, the compulsive hair removal behavior decreased and ultimately ceased following the dosage adjustment of her anxiolytic therapy, leading to the resolution of the skin manifestations. Considering the patient's clinical history, lesion characteristics, her reported compulsive use of tweezers on her face, and the improvement following optimization of anxiolytic therapy, we established a diagnosis of self-induced dermatitis factitia.

The pathological fiction of illness is found in all medical disciplines, with a variety of manifestations ranging from self-inflicted injuries to unnecessary interventions and repeated hospitalizations.² The patient's motivations and intentions, as well as the severity of manifestations and psychological correlates, vary widely. The ICD-11 currently defines artifactual skin disorders as a range of self-inflicted skin injuries that are provoked by mechan-




Figure 1. a, b) Skin lesions located on the left cheek, left eyebrow, and forehead that have been present for approximately two years; **c, d)** resolution of skin lesions after increasing the dosage of alprazolam for managing her anxious-depressive syndrome.

ical means or by the application or injection of chemical irritants or caustics.⁴ The true prevalence of these conditions is unknown, given their rarity and difficult diagnosis, but they are more frequently observed in females.⁵ Timely diagnosis of DF is essential to minimize the use of healthcare resources and mitigate potential harm to patients.⁴ Early intervention in the management of DF can also contribute to improved patient outcomes. The clinical and demographic characteristics of individuals affected by DF remain poorly defined.⁴ Most available literature on DF primarily comprises case reports and series, which, while important sources of information, may offer an incomplete clinical perspective of the disorder.³ A careful examination of skin lesions can raise suspicion, particularly when they exhibit atypical features.⁵ However, solely observing these lesions may not always be sufficient to suspect dermatitis factitia, as exemplified in our clinical case. Artefactual dermatitis should always be considered in the differential diagnosis of chronic, problematic, and recurrent skin disorders, particularly in patients who have psychiatric problems or specific psychological traits.⁶⁻⁸ When dermatitis presents with crusted lesions, it may be challenging to differentiate it from conditions like ecthyma and herpes simplex. In other instances, it can mimic photo-induced connectivitis, cutaneous sarcoidosis, vasculitis, and cutaneous lymphomas. In conclusion, DF continues to be a diagnosis of exclusion, necessitating appropriate investigations to rule out potential mimicking diseases.⁴ Apart from the imperative need for an early diagnosis, establishing a coordinated

treatment plan between the dermatologist and the psychiatrist is another crucial element in addressing FD effectively.-

References

1. Rook A, Wilkinson DS, Ebling FJB, et al. Textbook of dermatology. 4th ed. Oxford: Blackwell Scientific Publications; 1986.
2. Feldman MD, Eisendrath SJ. The spectrum of factitious disorders. *Am Psychiatr Publ* 1996.
3. Romano A, Alqahtani S, Griffith J, Koubeissi MZ. Factitious psychogenic nonepileptic paroxysmal episodes. *Epilepsy Behav Case Rep* 2014;2:184-5.
4. World Health Organization. ICD-11 revision. Available from: <https://icd.who.int/en>.
5. Mukundu Nagesh N, Barlow R, Mohandas P, et al. Dermatitis artefacta. *Clin Dermatol* 2023;41:10-5.
6. Wong JW, Nguyen TV, Koo JY. Primary psychiatric conditions: dermatitis artefacta, trichotillomania and neurotic excoriations. *Indian J Dermatol* 2013;58:44-8.
7. Yates GP, Feldman MD. Factitious disorder: a systematic review of 455 cases in the professional literature. *Gen Hosp Psychiatry* 2016;41:20-8.
8. Lavery MJ, Stull C, McCaw I, Anolik RB. Dermatitis artefacta. *Clin Dermatol* 2018;36:719-22.