

## **Dermatology Reports**

https://www.pagepress.org/journals/index.php/dr/index

eISSN 2036-7406







**Publisher's Disclaimer.** E-publishing ahead of print is increasingly important for the rapid dissemination of science. **Dermatology Reports** is, therefore, E-publishing PDF files of an early version of manuscripts that undergone a regular peer review and have been accepted for publication, but have not been through the copyediting, typesetting, pagination and proofreading processes, which may lead to differences between this version and the final one.

The final version of the manuscript will then appear on a regular issue of the journal. E-publishing of this PDF file has been approved by the authors.

Please cite this article as:

*Giorgio CM, Licata G, Tancredi V, et al.* Novel therapeutic approach for loxoscelism: efficacy of cefixime and netilmicin sulfate combination therapy. *Dermatol Rep 2025 [Epub Ahead of Print] doi: 10.4081/dr.2025.10215* 



Submitted 10/12/24 - Accepted 22/02/25

Note: The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

## Novel therapeutic approach for loxoscelism: efficacy of cefixime and netilmicin sulfate combination therapy

Caterina Mariarosaria Giorgio,<sup>1</sup> Gaetano Licata,<sup>2</sup> Vittorio Tancredi,<sup>1</sup> Paolino Franzese,<sup>1</sup> Nicola Maria Giorgio,<sup>1</sup> Giuseppe Argenziano,<sup>1</sup> Eugenia Veronica Di Brizzi<sup>1</sup>

<sup>1</sup>Dermatology Unit, Department of Mental and Physical Health and Preventive Medicine, University of Campania Luigi Vanvitelli, Naples; <sup>2</sup>Dermatology Unit, San Antonio Abate Hospital, Trapani, Italy

**Correspondence:** Gaetano Licata, MD, Dermatology Unit, San Antonio Abate Hospital, Via Cosenza 82, 91016, Erice (TP), Italy. Tel.: 3276215976 E-mail: gaetano.licata89@gmail.com

Key words: loxoscelism; cefixime; netilmicin sulfate; therapy.

Contributions: all authors equally contributed to data collection and article preparation.

Conflict of interest: the authors declare no potential conflict of interest.

**Ethics approval and consent to participate:** no ethical committee approval was required for this case report by the Department, because this article does not contain any studies with human participants or animals. Informed consent was obtained from the patients included in this study.

**Consent for publication:** the patients included in this manuscript have given written informed consent to the publication of their case details and any accompanying images.

## Dear Editor,

Loxoscelism, caused by bites from Loxosceles spiders, notably Loxosceles reclusa, presents with necrotic skin lesions and, in severe cases, systemic symptoms such as hemolysis, acute renal failure, and disseminated intravascular coagulation (DIC). The venom's sphingomyelinase D (SMD) enzyme induces direct tissue damage and an inflammatory response, leading to cell death, dermal necrosis, and increased vascular permeability, which can result in hemolysis and thrombosis.<sup>1,2</sup> Additionally, SMD activates endothelial cells and neutrophils, contributing to vasculitis, a factor often overlooked in treatment considerations. This mechanism may explain the efficacy of dapsone in certain stages of loxoscelism. In Italy, particularly in the southern regions, there has been an increase in reported Loxosceles spider bites over the past decade, correlating with climate changes and urban expansion into previously uninhabited areas.<sup>3</sup> This rise underscores the need for effective treatment protocols. Five patients presented to our clinic with confirmed Loxosceles spider bites, all exhibiting initial necrotic lesions (Figure 1 A,B). Previous treatments included local and systemic corticosteroids, topical antibiotics, and systemic antibiotics such as azithromycin and amoxicillin (administered separately), without significant improvement. The time between the spider bite and the initiation of these treatments ranged from 24 to 72 hours. Given the potential for secondary bacterial infections, we administered a combination of netilmicin (300 mg intramuscularly once daily) and cefixime (400 mg orally once daily) for 10 days. Netilmicin, an aminoglycoside antibiotic, is effective against a broad spectrum of gram-negative bacteria, including Pseudomonas aeruginosa,<sup>4</sup> and some grampositive bacteria. Cefixime, a third-generation oral cephalosporin, provides coverage against grampositive organisms like Streptococcus pyogenes and gram-negative bacteria, including Enterobacteriaceae.<sup>5</sup> Patients reported significant improvement in pain and local inflammation within the first five days of treatment (Figure 1C). All patients experienced resolution of their necrotic ulcers without the need for surgical intervention or further complications (Figure 1D). No side effects were reported. Wound cultures from two patients, taken before antibiotic administration, revealed the presence of Staphylococcus aureus and Pseudomonas aeruginosa, both of which were effectively cleared by the combination therapy. The combination of netilmicin and cefixime offers broadspectrum antibacterial coverage, addressing both gram-positive and gram-negative bacteria, including potential polymicrobial infections. This approach may be advantageous over conventional therapies that typically involve a single antibiotic or supportive wound care alone. However, the use of prophylactic antibiotics in loxoscelism should be carefully considered, as current guidelines do not universally recommend their use without clear signs of infection. In our cases, the presence of necrotic lesions and positive wound cultures justified the antibiotic therapy. The combination of netilmicin

and cefixime may provide an effective treatment for necrotic lesions resulting from *Loxosceles* spider bites, particularly when secondary bacterial infection is a concern. Further studies are warranted to establish the long-term efficacy and safety of this treatment protocol.

## References

1. Gremski LH, da Justa HC, Polli NLC, et al. Systemic Loxoscelism, Less Frequent but More Deadly: The Involvement of Phospholipases D in the Pathophysiology of Envenomation. Toxins (Basel) 2022;15:17.

2. Monteiro CL, Rubel R, Cogo LL, et al. Isolation and identification of Clostridium perfringens in the venom and fangs of Loxosceles intermedia (brown spider): enhancement of the dermonecrotic lesion in loxoscelism. Toxicon 2002;40:409-18.

3. Fagan SP, Berger DH, Rahwan K, Awad SS. Spider Bites Presenting with Methicillin-Resistant Staphylococcus aureus Soft Tissue Infection Require Early Aggressive Treatment. Surg Infect (Larchmt) 2003;4:311-5.

4. Caporossi A, Alessio G, Fasce F, et al. Short-Term Use of Dexamethasone/Netilmicin Fixed Combination in Controlling Ocular Inflammation After Uncomplicated Cataract Surgery. Clin Ophthalmol 2021;15:2847-54.

5. Arumugham VB, Gujarathi R, Cascella M. Third-Generation Cephalosporins. 2023 Jun 4. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024.

**Figure 1. A,B)** Initial clinical manifestation consisted of a beginning central necrotic area surrounded by erysipelas-like reaction. Lymphangitis is shown in panel A; **C)** after 5 days of combined antibiotic treatment, the erythema has significantly faded; **D)** complete resolution occurred after 10 days of treatment.

