

Dermatology Reports

https://www.pagepress.org/journals/index.php/dr/index

eISSN 2036-7406







Publisher's Disclaimer. E-publishing ahead of print is increasingly important for the rapid dissemination of science. **Dermatology Reports** is, therefore, E-publishing PDF files of an early version of manuscripts that undergone a regular peer review and have been accepted for publication, but have not been through the copyediting, typesetting, pagination and proofreading processes, which may lead to differences between this version and the final one. The final version of the manuscript will then appear on a regular issue of the journal. E-publishing of this PDF file has been approved by the authors.

Please cite this article as: Nsir S, Bouslama G, Ghanmi F, et al. The tongue: a bridge between physiology and disorder. *Dermatol Rep 2025 [Epub Ahead of Print] doi:* 10.4081/dr.2025.10262

@ the Author(s), 2025 *Licensee* <u>PAGEPress</u>, Italy

Submitted 16/01/25 - Accepted 27/01/25

Note: The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

The tongue: a bridge between physiology and disorder

Sabrine Nsir, Ghada Bouslama, Fadwa Ghanmi, Hanen Boukhris, Lamia Oualha, Souha Ben Youssef

Dentistry Department, Research Laboratory LR 12SP10: Functional and Aesthetic Rehabilitation of Maxillary, Farhat Hached Hospital, University of Sousse, Tunisia

Correspondence: Ghada Bouslama, Dentistry Department, Research Laboratory LR 12SP10: Functional and Aesthetic Rehabilitation of Maxillary, Farhat Hached Hospital, University of Sousse, 4000 Sousse, Tunisia.

E-mail: <u>bouslama.ghada@yahoo.fr</u> Tel.: +21622142003

Key words: tongue; pathology; depapillation.

Contributions: SN, GB, conceptualization, writing – original draft preparation; FG, methodology, writing – review & editing; HB, LO, investigation, writing – review & editing; SBY, formal analysis, writing – review & editing.

Conflict of interest: the authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Ethics approval and consent to participate: no ethical committee approval was required for this case report by the Department, because this article does not contain any studies with human participants or animals. Informed consent was obtained from the patients included in this study.

Consent for publication: informed consent for the publication of these review images was obtained from all patients.

Availability of data and materials: data sharing is not applicable to this article as no new data were created or analyzed in this study.

Abstract

The tongue, as a multifunctional organ, exhibits a variety of physiological characteristics as well as pathological alterations. Identifying and understanding these variations is crucial for ensuring an accurate diagnosis and appropriate management. This article explores the main common physiological variations, such as geographic tongue, fissured tongue, etc., as well as pathological conditions, including elementary lesions such as vesicles, bullae, erosions, ulcers, plaques, as well as surface changes (atrophy), color changes (coatings, pigmentation), or even shape alterations (infiltrations). It discusses current diagnostic tools and therapeutic approaches while emphasizing the importance of regular monitoring to prevent complications and maintain optimal oral health.

Introduction

The tongue is a multifunctional organ essential for speech, swallowing, and taste and also as an indicator of general health. Changes in color, appearance, and burning sensations can reveal a multitude of conditions, ranging from simple physiological variations to more serious pathologies.¹

Through the analysis of primary mucosal lesions such as vesicles, bullae, erosions, ulcers, plaques, changes in surface condition (atrophy), color (coatings, pigmentations), or even shape (infiltrations), an accurate diagnosis can be established.

Certain non-pathological aspects of the tongue may motivate a consultation and must be quickly recognized to reassure the patient of their benign nature.

This review article bridges the gap between normal physiological processes and pathological disorders, highlighting the tongue's importance as a window into overall health and offering insights into its potential in early disease detection.

Physiological variations of the tongue

Fissured tongue: a normal variant observed in 20 to 30% of the population, characterized by increased fissures and grooves on the dorsal and lateral surface (Figure S1).

It is a painless, most often isolated condition, though it can sometimes be associated with a geographical tongue. Along with facial edema (macrocheilitis in 66 to 96% of cases) and peripheral facial paralysis, it could be part of the diagnostic triad of Melkerson-Rosenthal syndrome. No specific treatment is necessary. Deeper fissures are often associated with a plicated tongue.^{2,3}

Geographic tongue: also known as benign migratory glossitis, it may be present at all ages. Clinically, it is characterized by areas of lingual depapillation, affecting the filiform papillae, delimiting erythematous regions with a slightly raised white border. These depapillated areas are irregularly arranged (Figures S2 and S3). Atypical forms may concern the ventral side or other extralingual oral regions called erythema migrans. No treatment is necessary for asymptomatic forms. An association between geographic tongue and cutaneous psoriasis is described with a common genetic determinism.¹⁻

Lingual varices: a common developmental abnormality in older adults. The etiology of oral varicosities remains unknown; they correspond to a benign vascular dilatation of the sublingual veins, non-inflammatory and painless, with degeneration of elastic fibers, which can be confused with an arteriovenous malformation (Figure S4). The diagnosis is clinical. No treatment is required.^{1,3}

Plica fimbriata: represents an embryonic remnant found under the tongue. These are folds of the mucosa that extend on either side of the lingual frenulum, forming slightly fringed structures (Figure S5).

Hypertrophy of circumvallate papillae, foliate papillae, and lingual tonsils: often a source of concern for some patients, which can sometimes simulate squamous cell carcinoma. The lingual tonsils are sometimes inflammatory and sensitive during infectious episodes of the upper aerodigestive tract (Figure S6).^{1,3}

Macroglossia: defined as an enlargement of the tongue that can lead to functional problems with ventilation and swallowing. It is characterized by the presence of tooth imprints on the lateral borders of the tongue and may even cause repetitive biting (Figure S7). Macroglossia can be due to congenital or acquired causes. Congenital factors include muscular hypertrophy, glandular hyperplasia, neonatal diabetes, and certain genetic syndromes such as Beckwith-Wiedemann syndrome, Melkersson-Rosenthal syndrome, and Down syndrome. Acquired factors include acromegaly, hypothyroidism, amyloidosis, and tongue tumors.³

Coatings

Black Hairy Tongue: refers to an elongation of the filiform papillae on the dorsal surface of the tongue, which are colored from brown to black (the tip of the tongue and the lateral surfaces are not affected). The coloration is due to the oxidation of keratin, which is attributed to the excessive use of antiseptic mouthwashes, tobacco, poor oral hygiene, or antibiotic therapy. Treatment involves eliminating risk factors and cleaning the dorsal surface of the tongue through scraping and brushing.^{1,3}

Saburral tongue: the dorsal surface of the tongue is covered with a whitish coating while the mucous membrane is normal below (Figure S8). This coating is made up of cellular desquamation, food debris, and saprophytic bacterial flora; it disappears when the cause is eliminated.^{1,3}

Pathological variations of the tongue

Pathological changes in the tongue can alter its color, texture, coating, and size, among other features. A comparison of normal and abnormal tongue characteristics is summarized in Table 1. Accurate clinical examination based on the analysis of primary lesions (ulceration, erosion, nodule, plaque, macule, papule, bullae, vesicles, etc.) is essential for distinguishing these variations.

Erythemas

Erythroplakia: a rare lesion with malignant potential, characterized by a velvety red area, well defined, and most often uniform without trace of keratinization. Risk factors include tobacco and alcohol. The rate of malignant transformation is much higher than that of leukoplakia,² making any erythroplakia lesion eligible to be biopsied and treated promptly. Long-term follow-up is necessary.⁴

Kawasaki disease: a vasculitis that tends to occur in children before the age of five. It is characterized by prolonged fever, bilateral conjunctivitis, exanthema, cervical lymphadenopathy, and palmar-plantar erythema with edema and oropharyngeal involvement. Oral involvement includes dry, cracked lips and the typical appearance of a strawberry tongue with prominent papillae.^{1,5}

Scarlet fever: a bacterial infection caused by group A *Streptococcus*, characterized by a rash, high fever, sore throat, and enanthema; the initially saburral tongue loses its whitish coating from the periphery to the center and becomes uniformly red with prominent papillae, also known as a strawberry tongue.¹

Anemia and vitamin deficiency: iron deficiency anemia is responsible for angular cheilitis with cracked lips and depapillation of the tongue. Treatment is based on anemia correction with iron supplementation.^{1,6} Anemia due to folate (vitamin B9) or vitamin B12 deficiency can cause Hunter's glossitis, which manifests itself at an early stage by lingual paresthesias, burning sensations, intolerance to dental prostheses, and/or dry mouth. Subsequently, red, glossy, depapillated areas are noticed.

The advanced phase of Hunter's glossitis shows a fissure network with a plicated appearance and can sometimes present a pultaceous coating (Figure S9). Etiological and replacement treatment of the deficiency allows the symptoms to disappear.^{1,7}

Oral candidiasis: can affect the entire oral mucosa. However, some variants can manifest exclusively on the tongue, such as median rhomboid glossitis. It is characterized by a rhomboid erythematous area on

the dorsal surface of the tongue, anterior to the lingual V. The area may be smooth and pink, flat, or papillary, often confused with hypertrophied circumvallate papillae. A mirror-image palatal erythematous lesion may be associated (Figures S10 and S11).⁸

The vesicles

Basic vesicular lesions are essentially linked to viral attacks, giving general and oral manifestations. These include herpes simplex virus (HSV) (Figure S12), varicella-zoster virus (VZV), and hand, foot, and mouth syndrome.¹ These viral infections generally result in painful post-vesicular erosions. However, some particularities are observed in the tongue, such as herpetic geometric glossitis, which often develops in immunocompromised patients. Lesions appear as branched longitudinal linear fissures, accompanied by extremely painful post-vesicular erosions preventing swallowing and speech.^{9,10} VZV can particularly affect the lingual nerve. In this case, the patient describes lingual pain, a decreased sensitivity, an alteration of taste, and eating discomfort associated with erosions covered with a pseudo-membranous coating complicating swallowing.

Bullae

Autoimmune bullous stomatitis, such as pemphigus, or cicatricial pemphigoid, or immuno-allergic stomatitis, such as erythema multiforme or hemorrhagic bullous angina, can all be revealed by lingual damage.¹ Erythema multiforme (EM) is characterized by targetoid skin lesions associated with lip involvement (Figure S13).¹¹ However, when it is accentuated (major EM), it presents lesions in the tongue and oral mucosa. Contrary to its name, hemorrhagic bullous angina can occur anywhere in the oral cavity, including the tongue, in the form of a large bulla with hematic content, giving way to painful ulceration.¹² Pemphigoid, in its cicatricial form (Figure S14), primarily affects the gingiva, presenting as erosive gingivitis. Lingual involvement is more frequent in pemphigoid compared to pemphigus vulgaris, which primarily affects the skin and presents as painful, irregularly shaped ulcers that hinder swallowing and speech. These conditions are often subject to candidal superinfections.¹³

Ulcers

Traumatic lesions: the tongue is more frequently affected by traumatic lesions due to its mobility. These lesions can generally give rise to ulcerations following physical trauma (Figure S15) (sharp edge, coronal fracture, hook, poorly adapted prosthesis), chemical (application of concentrated antiseptic, use of

tobacco) (Figure S16), and thermal (very hot foods or liquids). Removal of the causative agent allows the lesion to heal.¹⁴

Aphthous stomatitis and recurrent aphthous stomatitis: aphthous ulcers are very common lesions of the oral mucosa. They present as painful ulcers varying in size from a few millimeters (for minor aphthae) to up to two centimeters (for major aphthae), with a rounded or oval shape and surrounded by an erythematous halo (Figure S17). They typically occur on non-keratinized mucosa, including the lateral and ventral surfaces of the tongue, and rarely on keratinized mucosa (dorsal surface of the tongue). Minor aphthous ulcers heal without scarring, unlike major aphthae. Recurrent aphthous stomatitis is diagnosed if there are at least three episodes of multiple aphthous ulcers per year.¹⁴ Certain pathologies can cause aphthoid ulcerations, such as chronic inflammatory intestinal diseases, hematological disorders (anemia, neutropenia) (Figure S18), and taking certain medications (nicorandil, NSAIDs, bisphosphonate, etc.)

Eosinophilic ulcer: known as Riga-Fede disease. It is a rare benign lesion of the oral mucosa located mainly on the anterior part of the tongue following its friction on the edges of the temporary incisors during sucking. It presents as a painful ulcer, most often rounded, measuring a few mm to 2-3 cm in diameter, covered by a pseudo-membranous coating. Treatment is symptomatic with analgesic purposes.¹⁵

Ulcerations linked to vasculitis: Horton's disease: lingual damage can be a revealing sign of this arteritis: pain/burning, loss of taste sensitivity, claudication of the tongue associated with ulcerations rapidly evolving into necrosis and lingual gangrene.¹⁶

Bacterial origin: syphilis: an infectious sexually transmitted disease caused by *Treponema pallidum*. Primary syphilis, characterized by syphilitic chancre, can affect the entire oral mucosa without particularities in the tongue. On the other hand, in secondary syphilis, the lingual lesions are described as characteristic "mowed lawn" plaques. These damages can progress to necrotic ulcerations in the case of syphilitic gummas; we then speak of associated atrophic glossitis.¹

Malignant tumors: squamous cell carcinoma of the tongue: *is* the most common lymphophilic cancer, surpassing vertucous carcinoma and basal cell carcinoma. The most common sites are the ventral surface, lateral borders (Figure S19), and posterior border, which have a guarded prognosis compared to other locations with a delayed diagnosis (revealed with simple swallowing difficulty). The ulceration varies in size and has an irregular shape with an indurated, elevated, and sometimes everted edge. The base is indurated, extending beyond the ulceration and deep into the tissue.¹⁷

Nodules/papules

Generally testify to a benign attack most often linked either to an attack on the human papillomavirus (papilloma/wart) or a benign tumor such as fibroma, nerve tumors, which are most frequent in the tongue compared to other oral sites (neuroma, neurofibroma, schwannoma, Abrikossof tumor), and vascular tumors such as angioma, lymphangioma, and arteriovenous malformations, which can pose a risk of asphyxia.¹⁸

White lesions

Leukoplakia: a term used to clinically describe any white lesion of the oral mucosa idiopathic or associated with tobacco use. Lesions can be homogeneous or inhomogeneous, presenting a high risk of malignant transformation and requiring regular monitoring.²

Lichen planus: the tongue is the second oral site most frequently affected by this chronic inflammatory disease after the posterior jugal mucosa. Lingual lichen planus presents a particular form at the beginning, which manifests itself as more or less rounded plaques resembling "candle wax spots" (Figure S20). These lesions evolve inconsistently towards an erosive form under the influence of certain factors. In the long term, the mucosa becomes atrophic with the disappearance of the lingual papillae. This thin and fragile mucosa presents an increased risk of malignant transformation, requiring regular monitoring.¹⁴

Hairy oral leukoplakia: this lesion occurs on the borders of the tongue and is associated with Epstein-Barr virus, observed in immunocompromised individuals (HIV). Clinically, it presents as an asymptomatic white plaque with a hairy or undulating surface that resists scraping. However, rare cases affecting other sites of the oral mucosa have been observed.¹⁹

Uremic stomatitis: a rare condition in patients with renal failure. There are 4 forms of uremic stomatitis: ulcerative, hemorrhagic, pseudomembranous, and hyperkeratotic (Figure S21). The tongue and the buccal floor are most often affected. Associated symptoms are xerostomia, foul breath with an ammoniacal odor, bad taste, and burning sensations.¹

Thus, systemic diseases can often manifest with characteristic changes in the tongue, serving as valuable indicators in clinical diagnosis. Table 2 summarizes some common systemic conditions and their associated tongue manifestations.

Infiltrations

Amyloidosis: a rare disease characterized by the presence of amyloid deposits in various tissues and organs, including the oral cavity. Lingual damage can reveal macroglossia with indentations over the entire lingual margin (Figure S22). If the salivary glands are affected, xerostomia may develop. The

diagnosis is histological, showing positive Congo red staining, which indicates the presence of amyloid proteins. Amyloidosis is mostly found in patients with multiple myeloma.²

Others

Lingual abscess: more common in diabetic individuals, this condition often presents as a painful swelling that causes protrusion of the tongue, dysphagia, odynophagia, and difficulties with speech (Figure S23). Local trauma, such as injury, bite, spread of a dental infection, sting, or foreign body, is the primary predisposing factor for anterior abscesses. In contrast, lingual tonsillitis or an infected thyroglossal duct cyst are predisposing factors for posterior abscesses, which are considered potentially life-threatening. Management of a lingual abscess requires prompt and aggressive treatment: antibiotic therapy combined with surgical drainage.²⁰

Conclusions

Distinguishing between physiological and pathological variations of the tongue is crucial for accurate diagnosis and appropriate management. Although many variations are benign, it is essential to remain vigilant for abnormal changes in the tongue to prevent complications or detect undiagnosed systemic pathologies. Regular follow-up is recommended to maintain oral health.

References

1. Agbo-Godeau S, Weber P, Guedj A, Szpirglas H. "What can language examination reveal?".Current Odonto-Stomatol 2010;101-15.

2. Mangold AR, Torgerson RR, Rogers RS. Diseases of the tongue. Clin Dermatol 2016;34:458-69.

3. Vigarios E, De Bataille C, Boulanger M, et al. Physiological variations of the tongue. Ann Dermatol Venereology 2015;142:583-92.

4. Abati S, Bramati C, Bondi S, et al. Oral Cancer and Precancer: A Narrative Review on the Relevance of Early Diagnosis. Int J Environ Res Public Health 2020;17:9160.

5. Newburger JW, Takahashi M, Gerber MA, et al. Diagnosis, Treatment, and Long-Term Management of Kawasaki Disease. Traffic 2004;110:2747-71.

6. Seckinger C, Curien R, Hubert AC, et al. Oral manifestations of anemia. Current Odonto-Stomatol 2010;35-41.

7. Campana F, Sibaud V, Taieb A, Fricain JC. Oral manifestations revealing a vitamin B 12 deficiency: report of a case. Oral Med Oral Surg 2007;13:213-7.

8. Millsop JW, Fazel N. Oral candidiasis. Clin Dermatol 2016;34:487-94.

9. Mirowski GW, Goddard A. Herpetic geometric glossitis in an immunocompetent patient with pneumonia. J Am Acad Dermatol 2009;61:139-42.

10. Rocha-Méndez LE, Ochoa-Hein E, Gutiérrez-González AN, Tarango-Martínez VM. Herpetic geometric glossitis presenting as vegetative plaques in the context of acquired immunodeficiency syndrome. JAAD Case Rep 2021;12:64-6.

11. Rochefort J, Hervé G, Agbo-Godeau S. Oral manifestations of erythema multiforme: presentation of clinical cases. Oral Medicine Oral Surgery 2016;22:137-43.

12. Milin C, Fouche B. Hemorrhagic bullous angina after dental treatment: presentation of a case. Oral Medicine Oral Surgery 2011;17:293-6.

13. Nemnom PE, Baeck M, Dachelet C, Marot L. Autoimmune bullous diseases. French version available from: https://www.louvainmedical.be/sites/default/files/content/article/pdf/lm-11-2018-9-peause_dermato.pdf

14. Vigarios E, De Bataille C, Campana F, et al. Chronic or recurrent lingual ulcerations. Ann Dermatol Venereology 2016;143:297-308.

15. Samson J, El Hage M, Lombardi T. Don't always be fooled by age! Oral Med Oral Surg 2010;16:259-60.

16. Paraskevas KI, Boumpas DT, Vrentzos GE, Mikhailidis DP. Oral and ocular/orbital manifestations of temporal arteritis: a disease with deceptive clinical symptoms and devastating consequences. Clin Rheumatol 2007;26:1044-8.

17. Bilgic O, Duda L, Sánchez MD, Lewis JR. Feline Oral Squamous Cell Carcinoma: Clinical Manifestations and Literature Review. J Vet Dent 2015;32:30-40.

18. Müller S. Update from the 4th Edition of the World Health Organization of Head and Neck Tumors: Tumors of the Oral Cavity and Mobile Tongue. Head Neck Pathol 2017;11:33-40.

19. Alramadhan SA, Bhattacharyya I, Cohen DM, Islam MN. Oral Hairy Leukoplakia in Immunocompetent Patients Revisited with Literature Review. Head Neck Pathol 2021;15:989-93.

20. Srivanitchapoom C, Yata K. Lingual Abscess: Predisposing Factors, Pathophysiology, Clinical Manifestations, Diagnosis, and Management. Int J Otolaryngol 2018;2018:1-8.

 Table 1. Normal vs. abnormal tongue features.

Feature	Normal tongue	Abnormal tongue
Color	Pink, uniform color	White, red, pale, blue, purple or
		yellow: the tongue's color
		change could be a symptom of
		an underlying health condition
Texture	Smooth, slightly rough due to	Rough, swollen or smooth with
	papillae	a shiny appearance
Coating	Thin coating	Thick, yellow, white or black
		coating
Shape/Size	Normal	Enlarged, swollen or atrophic
Movability	Free, fluid, able to move in all	Restricted movement
	directions	
Edges	Normal, slightly tapered at the	Scalloped edges, indentations,
	tip	or teeth marks
Moisture	Moist, not excessively dry or	Dry or excessively moist tongue
	wet	

Table 2. Systemic diseases and their tongue manifestations.

Disease	Signs on the tongue	Clinical relevance
Anemia	Pale, smooth tongue with atrophy of papillae	associated with fatigue, weakness, and difficulty swallowing
Vitamin B12 Deficiency	Glossitis (inflamed, smooth, red tongue)	lingual paresthesias, burning sensations, intolerance to dental prostheses, and/or dry mouth
Diabetes	Dry, cracked tongue; may also show fungal infections (<i>e.g.</i> , candidiasis)	Infections and oral health issues: cavities/gum disease/periodontitis
Scarlet Fever	Strawberry tongue (red with prominent papillae)	Characterized by a rash, high fever, sore throat and enanthema
Horton's disease	Ulcerations rapidly evolving into necrosis and lingual gangrene	Pain/burning, loss of taste sensitivity and claudication of the tongue
Uremic stomatitis	4 forms: ulcerative, hemorrhagic, pseudomembranous and hyperkeratotic	Xerostomia, foul breath with an ammoniacal odor, bad taste and burning sensations

Online Supplementary Material