



eISSN 2036-7406

## Dermatology Reports

<https://www.pagepress.org/journals/index.php/dr/index>



**SIDCO**  
Società Italiana di Dermatologia  
Chirurgica, Oncologica, Correttiva ed Estetica

**Publisher's Disclaimer.** E-publishing ahead of print is increasingly important for the rapid dissemination of science. **Dermatology Reports** is, therefore, E-publishing PDF files of an early version of manuscripts that undergone a regular peer review and have been accepted for publication, but have not been through the copyediting, typesetting, pagination and proofreading processes, which may lead to differences between this version and the final one.

The final version of the manuscript will then appear on a regular issue of the journal.

E-publishing of this PDF file has been approved by the authors.

*Please cite this article as:*

*Tchernev G, Kordeva S. Dermatologic surgery rounds: double rotation (Yin-Yang) flap for reconstruction of a circular skin defect after BCC removal in the scalp region. Dermatol Rep 2025 [Epub Ahead of Print] doi: 10.4081/dr.2025.10420*

 © the Author(s), 2025  
Licensee [PAGEPress](https://www.pagepress.org/), Italy

Submitted 27/04/25 - Accepted 03/06/25

Note: The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

# **Dermatologic surgery rounds: double rotation (Yin-Yang) flap for reconstruction of a circular skin defect after BCC removal in the scalp region**

Georgi Tchernev,<sup>1,2</sup> Simona Kordeva<sup>1</sup>

<sup>1</sup>Onkoderma - Clinic for Dermatology, Venereology, and Dermatologic Surgery, Sofia; <sup>2</sup>Department of Dermatology and Venereology, Medical Institute of the Ministry of the Interior, Sofia, Bulgaria

**Correspondence:** Dr. Simona Kordeva, Onkoderma – Clinic for Dermatology, Venereology, and Dermatologic Surgery, General Skobelev 26, 1606 Sofia, Bulgaria.

E-mail: [simonakordeva97@gmail.com](mailto:simonakordeva97@gmail.com)

**Key words:** dermatologic surgery; double rotation flap; Yin-Yang flap; scalp; basal cell carcinoma.

**Conflict of interest:** the authors declare no potential conflict of interest.

**Ethics approval and consent to participate:** not required.

**Consent for publication:** the patient gave his written consent to use his personal data for the publication of this case report and any accompanying images.

## **The case**

An 84-year-old male came to the dermatology department with primary complaints of a tumor formation located on the vertex of the scalp. The surrounding skin showed mild sun damage, but no surgical scars or additional malignancies were noted. A clinical diagnosis of basal cell carcinoma was established, and surgical excision under local anesthesia with lidocaine was recommended.

## **Our choice**

The tumorous lesion located on the vertex of the scalp was surgically removed. The resultant primary defect was round, with exposed periosteum.

Medium-sized defects on the scalp, especially in hair-bearing areas, present a significant reconstructive challenge due to the size of the defect, the limited mobility of the scalp, and the sensitivity of its hair-bearing function. When selecting a reconstructive technique, it is essential to consider the scalp's mobility, the need for functional preservation, vascular integrity, tissue sparing, redistribution of tension vectors, and the desired aesthetic outcome. The proximity of several important anatomical structures must also be considered, including the supraorbital and supratrochlear nerves (branches of CN V1), the superficial temporal and posterior auricular arteries, and the superficial temporal vein.

We present a case of a primary medium-sized defect following surgical excision of a tumor located on the vertex of the scalp (Figure 1). Due to the size, location, and the involvement of a hair-bearing region, as well as the limited mobility and anatomical considerations of the scalp, our team decided against primary closure with single interrupted sutures or skin grafting. Potential risks such as functional impairment, aesthetic dissatisfaction, and increased tension on the wound edges were considered. Therefore, we opted for flap reconstruction using the patient's surrounding healthy tissue. This technique offers the advantages of maintaining similar anatomical integrity and preserving vascular supply while effectively reconstructing a medium-sized defect.

## **Procedure**

The lesion was surgically excised under local anesthesia with 2% lidocaine. Careful hemostasis was achieved. The resulting primary defect was reconstructed using the double rotation (Yin-Yang) flap technique (Figure 2). The rotation flaps were designed to be twice the length of the primary defect. Flap elevation was performed within the plane of the galea aponeurotica fascia to ensure vascular preservation. The flap vascularity relied on the superficial temporal and posterior auricular arteries. Dissection and undermining of the flaps down to the hypodermis were carefully performed, ensuring the preservation of the vascular supply. The pedicles consisted of subcutaneous tissue beneath the

flaps. Tension-free adaptation of the flaps was achieved, and the resulting secondary defect was closed with single interrupted 2-0 sutures (Figure 3a). The one-month postoperative period was uneventful, with no reported complications (Figure 3b).

### **Comment**

Scalp defect reconstruction poses a challenge for every dermato- or reconstructive surgeon due to the often larger wound defects, limited mobility of the scalp, the need to preserve function, protect the calvarium, and ensure the integrity of adjacent anatomical structures, often while aiming for an aesthetically acceptable outcome.<sup>1,2</sup> Primary wound closure in this area is usually preferred over secondary healing or skin grafting due to the limited applicability and increased risks associated with these two techniques in certain patients, such as those with a history of smoking, prior local radiation to the area, or where preservation of the scalp's hair-bearing function is needed.<sup>2,3</sup>

A thorough understanding of surgical anatomy, defect size, individual patient considerations, reconstructive goals, and realistic expectations is essential when selecting the most appropriate reconstructive technique.<sup>3</sup> Various reconstructive techniques, including advancement, rotation, and transposition flaps, can be employed to help distribute the tension vectors generated by the defect.<sup>1,3</sup> Challenging anatomical sites such as the scalp can be effectively managed using the bilateral rotation (Ying-Yang) flap, particularly when tension at the incision site is a concern.<sup>4</sup> The Ying-Yang flap is a double-opposing rotation advancement flap that allows for the redistribution of tension across multiple vectors.<sup>4</sup> Although not a novel approach, this method remains a safe, reliable, and effective reconstructive technique for managing primary defects of the scalp.

### **The outcome**

The outcome is shown in Figure 3 a,b.

### **References**

1. Talevi D, Torresetti M, Recchi V, Di Benedetto G. Moving from the O-Z flap to the O-S flap for scalp reconstruction: A new geometrical model. *JPRAS Open* 2024;42:178-85.
2. Bradford BD, Lee JW. Reconstruction of the Forehead and Scalp. *Facial Plast Surg Clin North Am* 2019;27:85-94.
3. Desai SC, Sand JP, Sharon JD, et al. Scalp reconstruction: an algorithmic approach and systematic review. *JAMA Facial Plast Surg* 2015;17:56-66.
4. Suárez Ortega HM, Ortega Landeros JJ, Mendoza Argáez JI, et al. Scalp Spinal Cell Carcinoma Managed with Resection and Ying Yang Flap. *Int J Med Sci Clin Res Stud* 2024;4:807-9.



**Figure 1.** A rounded tumorous formation located in the vertex region of the scalp, prominent above the surrounding skin, with ulcerated surface and covered in places with hemorrhagic crusts.



**Figure 2.** Intraoperative view: primary wound defect reconstructed using the double rotation (Yin-Yang) flap technique. The rotation flaps are designed to be twice the length of the primary defect. Flap elevation is performed within the plane of the galea aponeurotica fascia.



**Figure 3. a)** Intraoperative view: the secondary wound defect is closed with single interrupted sutures; **b)** one-month postoperative period.

