



Title: OSTEONECROSIS OF JAW (ONJ): IMPACT OF ITALIAN PATIENTS, AND
ROLE OF ITALIAN PHYSICIANS, DENTISTS, AND RESEARCHERS IN THE
GROWING EVIDENCE OF A "NEW" DISEASE

Authors: Fusco V.;^{1*} Baraldi A.;² Fasciolo A.;³ Rapetti M.;⁴ De Martino I.;³ Randi L.;³
Bedogni A.;⁵ Agrillo A.;⁶ Campisi G.;⁷

Type: Original Article

Keywords: Osteonecrosis of jaw;

Abstract

Purpose: Osteonecrosis of jaw (ONJ) is an uncommon but severe complication observed mostly in patients treated with bisphosphonates (BPs) for bone metastases, myeloma, osteoporosis (so called BRONJ, Bisphosphonate-Related Osteonecrosis of Jaw), but also with other drugs (bevacizumab, sunitinib, denosumab). The number of cases observed in Italy appears high in comparison with other countries and we present a review of several aspects of ONJ in Italy and the role of Italian health professionals and researchers on increasing knowledge and adequate reporting of ONJ phenomenon;

¹ Unit of Oncology, Department of Oncology and Haematology,

* Corresponding author: vfusco@ospedale.al.it;

² Unit of Haematology, Department of Oncology and Haematology,

³ Unit of Maxillo-Facial Surgery;

⁴ Centro Documentazione e Prevenzione Osteonecrosi,
Azienda Ospedaliera di Alessandria (City Hospital), Alessandria, Italy

⁵ Section of Oral and Maxillofacial Surgery, Department of Surgery,
Azienda Ospedaliera Universitaria Integrata, Verona, Italy

⁶ Section of Maxillofacial Surgery, Department of Odontostomatologic Sciences,
University "La Sapienza", Rome, Italy

⁷ Sector of Oral Medicine "V.Margiotta", Department of Surgical and Oncological Sciences,
University of Palermo, Palermo, Italy



Methods: Literature review about osteonecrosis of jaw (ONJ) with selection of Italian authors and publications, on year 2003-2011, by research on international electronic journal databases, Italian language journals, congress acta, web sources;

Results: at October 2011, among 1272 papers published worldwide on ONJ issue, 128 (10%) were from Italian Authors;

Conclusions: relevant articles by Italian groups were published about pathogenesis hypotheses, animal models, biology studies, risk factors, preventive measures, dental extraction protocols in BP-exposed patients, laser therapy, ozone therapy, surgical treatment. Experience of Italian patients suffering from ONJ, together with work of Italian dentists, physicians and researchers, appears of paramount importance in order to study ONJ and minimize a possible severe side-effect of efficacious medical treatments.

1. Introduction

Osteonecrosis of jaws (ONJ) has been largely described since first 2000 years [1-4] in patients receiving bisphosphonates (BPs), drugs widely administered for the treatment of bone disease in multiple myeloma and metastatic solid tumors but also in osteoporosis and other non-malignant bone pathologies. The mechanisms underlying the development of Bisphosphonate-Related ONJ (BRONJ) are not completely understood [5-6]. Furtherly, ONJ has been recently observed also in patients treated with other drugs, such as bevacizumab, sunitinib, and denosumab [7-9]. In Italy, the number of BRONJ cases observed is particularly high in comparison with other countries [10]. We present a review of several aspects of ONJ in Italy and the role of Italian health professionals and researchers on increasing knowledge and adequate reporting of ONJ phenomenon.

2. Material and methods

A manual searching was performed using a full text electronic journal database (Pub Med); the main applied key was "Osteonecrosis AND (jaw OR jaws)", and publications including Italian authors were selected.



Further research was conducted on Pub Med using terms : "Diphosphonates" [Mesh] OR Bisphosphonate* OR Diphosphonates*", "Breast Neoplasms" [Mesh], "Prostatic Neoplasms" [Mesh], " Multiple myeloma " [Mesh] AND "osteonecrosis".

Italian language journals (of oncology, haematology, maxillofacial surgery and dentist interest) not included on Pub Med were reviewed, since 2003 to October 2011.

National and international congress and conference acta (abstract books, educational books, highlights, etc.), both in English and Italian language, were examined to search possible contributions of interest.

Reports from AIFA (the Italian Drug Surveillance Agency) including ONJ cases and recommendations were considered. Guidelines from several Italian scientific societies and deliberations from Italian Ministry of Health about prevention and early diagnosis of ONJ were examined.

Further papers, recommendations, position papers, books, documents were found by a manual searching on the web.

3. Results

Pub Med publications

Between 2003 and October 2011, a large amount of articles have been published worldwide on ONJ issue (1272); 10 % of them (128) were from Italian Authors. We classified the latter items as follows: 38% are case reports and clinical experiences; 13 % report a therapeutic or behaviour protocol; 32% were about therapy; 13% were guidelines about preventive measures; 15% were literature reviews.

The analysis showed that 55% of publications by Italian authors were issued in the last three years (2009-2011), representing the increased interest and recent work of Italian physicians and researchers towards ONJ.

BRONJ pathogenesis hypotheses, animal models, pathological and genetics studies were objects of papers by several groups, located in Verona [11-12], Trieste [13] , Torino [14-15] , Florence [16], Bari [17] , Messina [18-21], and other academic centers [22-24].

Several BRONJ case reports and case series in cancer and myeloma patients have been published since first years of BRONJ appearance [25-40]. Particular case reports included one case of BRONJ associated to osteonecrosis of left thumb [41], BRONJ diagnosis years



after BP discontinuation [42], BRONJ in a hemodialysis patient [43], ONJ in a leukaemia patient not treated with BPs [44], BRONJ associated with photodynamic therapy [45].

BRONJ incidence (or prevalence, or frequency) data have been evaluated and discussed by Italian authors : in breast cancer patients [27,33,46-47], in prostate cancer patients [48-49] , in myeloma patients [50-54] , and in mixed neoplasms patient series [54-57] .

Large (more than 100 BRONJ cases) single-centre experiences have been published, such as that of Parma [58] and Catania [59] , or are under reviewing (e.g. Verona, Palermo, Rome, Milan, etc).

One regional experience (247 cases in Piemonte-Valle d'Aosta area, on years 2005-2008) has been published only as a summary, at this date [56, 60] .

A multicenter case collection of 672 BRONJ cases observed in several Italian centers, collected on 2009, is under evaluation and has generated one published article [61] and other papers under authors' discussion or under submission.

ONJ definition and staging were examined by several Italian researchers, who underlined problems concerning cases without bone exposure and the so-called "stage 0 " according to the AAOMS staging system [62-63] ; a proposal for a new staging system is under discussion by members of a panel prompted by Italian societies of maxillofacial surgeons (SICMF) and oral care specialists (SIPMO).

Imaging studies have been performed [64-69] and others are submitted or ongoing.

Risk factors are fundamental to know odd ratio of single patients. Systemic risk factors, such as type of BP (with zoledronic acid linked to higher risk) [46, 53] and treatment duration and schedule [53,70-72] have been studied; after-treatment residual risk was reported [42]. Local (dental) risk factors, including tooth extractions [70] , periodontal disease [25] , and dental implant [73-75] have been underlined.

About preventive measures, Italian authors published proposals of risk management [76-84] but also efficacy studies [85-86] and general population evaluations [60].

Dental extraction protocols in BP-exposed patients have been reported by several groups [12, 75, 87-89] in order to minimize the risk of ONJ after dentoalveolar trauma.

BRONJ in non-malignant diseases, mostly osteoporosis patients, was critically evaluated [80,82, 90-91] , with increased reports in latest years [73, 92-94]; Italian authors were included in an international multi-centre study [95] .



Reports and evaluations by Italian authors about ONJ in patients receiving other drugs (with or without BPs) have been published; they included bevacizumab [47, 96], sunitinib [97-98], denosumab [10] .

Uncertainty of efficient BRONJ treatment options was examined by several Italian groups [39, 55, 61, 99- 101]. Conservative treatments were indicated as fundamental by some authors [59, 93, 102-103] . Surgical treatment series or case reports have been object of several papers [61,104-111] , including a prospective experience of large jaw resections with long-term follow-up [112] .

The first case of autologous bone marrow stem cell intralesional transplantation to repair BRONJ has been recently published by an Italian centre [113] .

Laser therapy, with different instruments, was proposed by several groups [114-119] .

Also ozone therapy has been applied in different forms [120-123] , and comparison studies are needed to confirm its usefulness.

Italian language publications

In years 2004-2011, at least 84 Italian language articles (reviews, case reports, case series) and abstracts have been published (mostly in dentist Italian literature, but often on websites or web journals), even from physicians and centres not publishing on Pub Med journals .

Two Italian language books [124-125] and the translation of a textbook by RE Marx [126] have been printed.

AIFA reports

Since 2005, numerous reports from AIFA (the Italian Drug Surveillance Agency) included ONJ case series and prevention recommendations. However only a little part of ONJ cases published in literature and/or reported at congresses and conferences are also signalled to the Agency, due to a poor attitude of Italian physicians and dentists to centralized drug side-effect reporting (only 425 BRONJ cases had been reported to the national Drug Surveillance System as of June 2009) [127].

The AIFA offices actively worked with European Agency (EMEA, now EMA) on discussing ONJ phenomenon [128], both related to BP treatment and to other drugs (antiangiogenic agents) [129], collaborating on releasing of bevacizumab and sunitinib alerts on autumn 2010 [130, 131, 132] .



Italian Congresses

Beside capillary local/regional dentist conferences (mostly organized by ANDI, the Italian National Dentist Association) and local medical oncology/haematology symposia, we found ONJ-dedicated sessions or presentations in national congresses of several societies. Italian societies of maxillofacial surgeons (SICMF), oral pathologists and care specialists (SIPMO), oncologists (AIOM), haematologists (SIE) organized oral presentations, issue symposia and abstract discussion sessions in their congresses in latest years. Furtherly, dozens of abstracts and posters about BRONJ have been offered by scholars and students at annual meetings of the Italian college of oral care teachers ("Collegio dei Docenti di Odontoiatria").

Moreover, some specific-issue national conferences (Alessandria 2007, Messina 2008, Alessandria 2008-2009-2010) were organized, with dozens of abstracts and reports from centers of all the Italian regions (only minimally later published as full papers). Highlights of those national congresses have been diffused by Italian language newsletters [133] and most of contributions are available on a website (www.reteoncologica.it) [134].

European and International Congresses

Abstracts from Italian Authors about ONJ appeared in the acta of many international society congresses, such as ASH (American Society of Haematology), ASCO (American Society of Clinical Oncology), EHA (European Haematology Association), ESMO (European Society of Medical Oncology), ECCO (European Cancer Organization), EAOM (European Oral Medicine Association), EACMFS (European Association of Maxillo-Facial Surgeons), and international conferences (for example SABCS, San Antonio annual breast cancer symposium).

Recommendations, guidelines, position papers

Beside an official recommendation by Italian Health Minister [135], documents about ONJ were approved and diffused (or are going to be published) by several Italian societies and study groups: SICMF-SIPMO panel (an expert panel organized by Italian societies of maxillofacial surgeons and oral medicine specialists), AIOM (oncologists), SIF (pharmacologists), SIOMMMS (bone disease specialists), GISBI (BP study group), PROOF (dental legal experts), Rete Oncologica Piemonte-VdA (multidisciplinary study group about BP prescription in cancer and myeloma), etc.



Epidemiologic data

Systematic BRONJ case collection has been pursued on two regional area.

In Piemonte – Valle d’Aosta (4.3 million population), inside a regional oncology network (Rete Oncologica Piemonte Valle d’Aosta), on November 2005 a BRONJ multidisciplinary study group was created, after observation of 60 cases in few referral centers [136]; enlargement of the group to 37 oncology / haematology regional units and 19 oral care units pushed a systemic collection up to 247 cases on December 2008 [60] .

In Sicily, a local drug side-effect surveillance system (hospital pharmacist network) has collected 102 cases in 3 referral centers of two cities (Palermo and Messina) on years 2008-2009 [137] and is enlarging to all the region.

4. Discussion

In recent years many Italian groups have published a large amount of articles about several aspects of ONJ in English language peer-reviewed journals, reporting an high number of observed patients. An even larger amount of ONJ cases (mostly BRONJ in cancer and myeloma patients, but with increasing number in not malignant diseases) has been illustrated in national and international congresses and conferences. We calculated that Italian authors have reported between 2005 and 2009, either in congresses, dedicated annual meetings [133-134] and/or peer-reviewed journals, at least 1,200 different ONJ cases (among which are 672 cases from a multicenter survey, whose results are only partially published) [61]. Everyone could imagine that many other cases have not been reported at all.

Projecting to whole Italian population (58 million) the number of BRONJ cases (247) observed in Piemonte-Valle d’Aosta (4.3 million) in years 2004-2008 [60] , one could speculate that more than 3000 BRONJ cases have occurred in Italy in those years.

Why so many BRONJ cases in Italy, especially among cancer and myeloma patients? Looking for the main reasons, we invite readers to consider several possible facts. Firstly, on the basis of prescriptions in years 2002-2008 of zoledronic acid (the mostly administered iv BP, largely replacing pamidronate) (fig 1), we can consider that iv BP consumption rate in metastatic bone and myeloma patients in recent years was higher in Italy than in other European countries (with exception of Belgium) (AIFA personal communication) (fig 2); drug therapy is largely covered by the national public Italian Health System (SSN, Sistema Sanitario Nazionale), and this is totally free for oncology and haematology patients , in years

2001-2006 largely treated with BPs in public and private centres, on a monthly basis, accordingly to less recent guidelines indicating continuous BP treatment “*until evidence of substantial decline in a patient's general performance status*” [138-140] . Secondly, on the contrary, the SSN did not adopt adequate coverage for oral health measures, that are mainly left to individual choice and are *mostly* under care of private dentists; direct and indirect measures indicate an insufficient median status of oral health in Italian people, especially of middle-age and elderly people (the age in which you can find most of cancer and myeloma patients). Consequently, the sum of systemic factors (large amounts of iv monthly BPs, for prolonged time, particularly of zoledronic acid) and of local factors (bad oral health status; periodontal disease; high number of teeth needed to be extracted without specific protocol during BP treatment; etc) could explain such a high number of BRONJ cases in years 2004-2008.

In Italy diffusion of knowledge of BRONJ among specialists and medical and dentist practitioners was late in comparison with North America. In USA, since 2004 Novartis Pharmaceuticals Inc , the major manufacturer of large-diffused iv bisphosphonates, pamidronate (Aredia®) and zoledronic acid (Zometa®) alerted involved specialists (oncologists, haematologists, dentists, maxillofacial surgeons) with a “white paper” drawn up in March 2004 and made public in June 2004, followed by a “dear doctor” letter (September 2004); patients were even alerted through meetings with patient advocacy groups since May 2004 and the diffusion of a patient information booklet [4]. In Italy , only on late 2005 and on 2006, after stimulation [141] and after first publications on main international medicine journals [142], BP manufacturers and health authorities adopted measures to inform involved specialists. Afterwards, especially since 2007, information about this “new disease” was diffused, also among dentist practitioners. However, also in Italy [143] as well as in other countries [144-145] patients ignorance about administered BP treatment , as well as dentists and physicians insufficient information, remains a substantial risk factor for inducing new cases of BRONJ .

On the other side, the reduction of number of new BRONJ cases in latest years in some large Italian referral centers (data not yet published) as well in one regional area [60] seem indicate encouraging results from adoption of preventive measures, together with a more selective use of BPs (as indicated by more recent guidelines) [146-147] .

In conclusion: experience of Italian patients suffering from ONJ, together with work of Italian dentists, physicians and researchers, appears of paramount importance in order to



study ONJ and minimize as much as possible this severe side-effect of efficacious medical treatments.

References

- [1] R.E. Marx, "Pamidronate (Aredia) and Zoledronate (Zometa) induced avascular necrosis of the jaws: a growing epidemic," *J Oral Maxillofac Surg.*, vol. 61, pp. 1115-1118, 2003.
- [2] S.L. Ruggiero, J. R. Tracey, S.L. Engroff "Osteonecrosis of the Jaws Associated With the Use of Bisphosphonates: A Review of 63 Cases," *J Oral Maxillofac Surg.*, vol.62 pp.527-534, 2004.
1. [3]C.A. Migliorati "Bisphosphonates and oral cavity avascular bone necrosis". *J Clin Oncol.* Vol. 21(22): pp 4253-4, 2004.
2. [4] S.L. Ruggiero , "Bisphosphonate-related osteonecrosis of the jaw (BRONJ): initial discovery and subsequent development" *J Oral Maxillofac Surg.*, Vol. 67 (suppl 5), pp.13-18, 2009.
3. [5] M.R. Allen, D.B. Burr," The Pathogenesis of Bisphosphonate-Related Osteonecrosis of the Jaw: So Many Hypotheses, So Few Data," *J Oral Maxillofac Surg.*, vol. 67, pp. 61-70, 2009.
- [6] C.A. Migliorati, J.B. Epstein, E. Abt, J.R. Berenson, "Bisphosphonate-associated osteonecrosis (BON) is a complication," *Nat. Rev. Endocrinol.*, vol. 7, pp. 34-42, 2011.
- [7] C.L. Estilo, M. Fornier, A. Farooki, D. Carlson, G. Bohle III, J.M. Huryn, "Osteonecrosis of the Jaw Related to Bevacizumab," *Journal of Clinical Oncology*, vol. 26, no. 24, pp. 4037-4043, 2008.
- [8] J. Ayllon, V. Launay-Vacher, J. Medioni, C. Cros, J. P. Spano, S. Oudard, "Osteonecrosis of the jaw under bisphosphonate and antiangiogenic therapies: cumulative toxicity profile?"*Annals of Oncology* 20(3): pp 600-1, 2009.
- [9] T. L. Aghaloo, A. L. Felsenfeld, S. Tetradis, "Osteonecrosis of the Jaw in a Patient on Denosumab," *J Oral Maxillofac Surg.*, 68(5): pp 959-63, 2010.
- [10] V. Fusco, C. Galassi, A. Berruti, L. Ciuffreda, C. Ortega, G. Ciccone, A. Angeli, O. Bertetto, "Osteonecrosis of the jaw after zoledronic acid and denosumab treatment," *J Clin Oncol.*, vol. 29, no. 17, pp. 521-522, 2011.
- [11] F. Bertoldo, D. Santini, V. Lo Cascio, "Bisphosphonates and osteomyelitis of the jaw: a pathogenic puzzle," *Nat Clin Pract Oncol.*, vol. 4, no. 12, pp. 711-721, 2007.



- [12] G. Saia, S. Blandamura, G. Bettini, A. Tronchet, A. Totola, G. Bedogni, G. Ferronato, P. F. Nocini, A. Bedogni, "Occurrence of Bisphosphonate-Related Osteonecrosis of the Jaw After Surgical Tooth Extraction," *Oral Maxillofac Surg.*, vol. 68 pp.797-804, 2010.
- [13] M. Biasotto, S. Chiandussi, S. Zacchigna, S. Moimas, F. Dore, G. Pozzato, F. Cavalli, F. Zanconati, L. Contardo, M. Giacca, R. Di Lenarda, "A novel animal model to study non-spontaneous bisphosphonates osteonecrosis of jaw," *J Oral Pathol Med.*, vol. 39, no. 5, pp. 390-396, 2010.
- [14] M. Ardine, D. Generali, M. Donadio, S. Bonardi, M. Scoletta, A.M. Vandone, M. Mozzati, O. Bertetto, A. Bottini, L. Dogliotti, A. Berruti. Could the long-term persistence of low serum calcium levels and high serum parathyroid hormone levels during bisphosphonate treatment predispose metastatic breast cancer patients to undergo osteonecrosis of the jaw? *Ann Oncol.* Aug;17(8): pp 1336-7, 2006.
- [15] P.G. Arduino, E. Menegatti, M. Scoletta, C. Battaglio, M. Mozzati, A. Chiecchio, D. Berardi, A.M. Vandone, M. Donadio, S. Gandolfo, C. Scully, R. Broccoletti, "Vascular endothelial growth factor genetic polymorphisms and haplotypes in female patients with bisphosphonate-related osteonecrosis of the jaws," *J Oral Pathol Med*, vol. 40, no. 6, pp. 510-515, 2011.
- [16] F. Marini, P. Tonelli, L. Cavalli, T. Cavalli, L. Masi, A. Falchetti, M.L. Brandi, "Pharmacogenetics of bisphosphonate-associated osteonecrosis of the jaw," *Front Biosci (Elite Ed)*, vol. 1, no. 3, pp. 364-370, 2011.
- [17] G. Favia, G.P. Pilolli, E. Maiorano, "Histologic and histomorphometric features of bisphosphonate-related osteonecrosis of the jaws: an analysis of 31 cases with confocal laser scanning microscopy," *Bone*, vol. 45, no. 3, pp. 406-413, 2009.
- [18] A. Alonci, A. Allegra, G. Bellomo, E. Quartarone, G. Oteri, E. Nastro, D. Cicciù, F.S. De Ponte, C. Musolino, "Patients with bisphosphonate-associated osteonecrosis of the jaw have unmodified levels of soluble vascular endothelial growth factor receptor 1," *Leuk Lymphoma.*, vol. 48, no. 9, pp. 1852-1854, 2007.
- [19] G. Oteri, A. Allegra, G. Bellomo, A. Alonci, E. Nastro, G. Penna, L. Catalfamo, D. Cicciù, F.S. De Ponte, C. Musolino, "Reduced serum levels of Interleukin 17 in patients with osteonecrosis of the jaw and in multiple myeloma subjects after bisphosphonates administration," *Cytokine.*, vol. 43, no. 2, pp. 103-104, 2008.
- [20] A. Allegra, A. Alonci, G. Penna, A. Granata, E. Nastro Siniscalchi, G. Oteri, S. Loddo, D. Teti, D. Cicciù, F.S. De Ponte, C. Musolino, "Bisphosphonates induce apoptosis of



circulating endothelial cells in multiple myeloma patients and in subjects with bisphosphonate-induced osteonecrosis of the jaws," *Acta Haematol.*, vol. 124, no. 2, pp. 79-85, 2010.

[21] E. Nastro Siniscalchi, G. Cutroneo, L. Catalfamo, G. Santoro, A. Allegra, G. Oteri, D. Ciccì, A. Alonci, G. Penna, C. Musolino, F. Saverio De Ponte, G. Anastasi, A. Favalaro, "Immunohistochemical evaluation of sarcoglycans and integrins in gingival epithelium of multiple myeloma patients with bisphosphonate-induced osteonecrosis of the jaw," *Oncol Rep.*, vol. 24, no. 1, pp. 129-134, 2010.

[22] D. Baldi, A. Izzotti, P. Bonica, P. Pera, A. Pulliero, "Degenerative periodontal-diseases and oral osteonecrosis: the role of gene-environment interactions," *Mutat Res.*, vol. 667, no. 1-2, pp. 118-131, 2009.

[23] I. Perrotta, M.G. Cristofaro, M. Amantea, E. Russo, S. De Fazio, V. Zuccalà, F. Conforti, A. Amorosi, G. Donato, S. Tripepi, M. Giudice, "Jaw osteonecrosis in patients treated with bisphosphonates: an ultrastructural study," *Ultrastruct Pathol.*, vol. 34, no. 4, pp. 207-213, 2010.

[24] M.T. Di Martino, M. Arbitrio, P.H. Guzzi, E. Leone, F. Baudi, E. Piro, T. Prantera, I. Cucinotto, T. Calimeri, M. Rossi, P. Veltri, M. Cannataro, P. Tagliaferri, P. Tassone, "A peroxisome proliferator-activated receptor gamma (PPARG) polymorphism is associated with zoledronic acid-related osteonecrosis of the jaw in multiple myeloma patients: analysis by DMET microarray profiling," *Br J Haematol.*, vol. 154, no. 4, pp. 529-533, 2011.

[25] G. Ficarra, F. Beninati, I. Rubino, A. Vannucchi, G. Longo, P. Tonelli, G. Pini Prato, "Osteonecrosis of the jaws in periodontal patients with a history of bisphosphonates treatment," *J Clin Periodontol.*, vol. 32, no. 11, pp. 1123-1128, 2005.

[26] A.M. Vannucchi, G. Ficarra, E. Antonioli, A. Bosi, "Osteonecrosis of the jaw associated with zoledronate therapy in a patient with multiple myeloma," *Br J Haematol.*, vol. 128, no. 6, pp. 738, 2005.

[27] V. Guarneri, S. Donati, M. Nicolini, S. Giovannelli, R. D'Amico, P.F. Conte, "Renal safety and efficacy of i.v. bisphosphonates in patients with skeletal metastases treated for up to 10 Years," *Oncologist.*, vol. 10, no. 10, pp. 842-848, 2005.

[28] A. Oltolina, A. Achilli, G. Lodi, F. Demarosi, A. Sardella, "Osteonecrosis of the jaws in patients treated with bisphosphonates. Review of the literature and the Milan experience," *Minerva Stomatol.*, vol. 54, no. 7-8, pp. 441-448, 2005.



- [29] E. Merigo, M. Manfredi, M. Meleti, D. Corradi, P. Vescovi, "Jaw bone necrosis without previous dental extractions associated with the use of bisphosphonates (pamidronate and zoledronate): a four-case report," *Oral Pathol Med.*, vol. 34, pp. 613-617, 2005.
- [30] M.D. Mignogna, S. Fedele, L. Lo Russo, R. Ciccarelli, L. Lo Muzio, "Case 2. Osteonecrosis of the jaws associated with bisphosphonate therapy," *J Clin Oncol.*, vol. 24, no. 17, pp. 2687, 2006.
- [31] R. Bertè, A. Arcari, P. Bernuzzi, E. Anselmi, A. Lazzaro, C.F. Moroni, D. Vallisa, L. Cavanna, "Jaw avascular bone necrosis associated with long-term use of bisphosphonates," *Tumori*, vol. 92, no. 4, pp. 361, 2006.
- [32] C. Ortega, R. Faggiuolo, R. Vormola, F. Montemurro, D. Nanni, F. Goia, M. Aglietta, "Jaw complications in breast and prostate cancer patients treated with zoledronic acid," *Acta Oncol.*, vol. 45, no. 2, pp. 216-217, 2006.
- [33] G. Sanna, L. Preda, R. Bruschini, M. Cossu Rocca, S. Ferretti, L. Adamoli, E. Verri, L. Franceschelli, A. Goldhirsch, F. Nolè, "Bisphosphonates and jaw osteonecrosis in patients with advanced breast cancer," *Ann Oncol.*, vol. 17, no. 10, pp. 1512-1516, 2006.
- [34] D. Errante, N. Mannucci, D. Bernardi, A. Bianco, L. Salvagno, "Comment on 'Jaw avascular bone necrosis associated with long-term use of bisphosphonates'," *Ann Oncol.*, vol. 17, no. 2, pp. 350-352, 2005.
- [35] N. Salesi, R. Pistilli, V. Marcelli, F.A. Govoni, F. Bozza, G. Bossone, V. Venturelli, B. Di Cocco, U. Pacetti, A. Ciorra, C. Di Fonso, E. Cortesi, E. Veltri, A. Vecchione, "Bisphosphonates and oral cavity avascular bone necrosis: a review of twelve cases," *Anticancer Res.*, vol. 26, no. 4B, pp. 3111-3115, 2006.
- [36] E. Merigo, M. Manfredi, M. Meleti, R. Guidotti, A. Ripasarti, E. Zanzucchi, P. D'Aleo, D. Corradi, L. Corcione, E. Sesenna, S. Ferrari, T. Poli, M. Bonanini, P. Vescovi, "Bone necrosis of the jaws associated with bisphosphonate treatment: a report of twenty-nine cases," *Acta Biomed.*, vol. 77, no. 2, pp. 109-117, 2006.
- [37] S. Capalbo, M. Delia, D. Diomede, M. Dargenio, A. Chiefa, G. Favia, V. Liso, "Jaw osteonecrosis associated with use of bisphosphonates and chemotherapy: paradoxical complication of treatment of bone lesions in multiple myeloma patients," *Int J Hematol.*, vol. 83, no. 5, pp. 439-442, 2006.
- [38] C. Broglia, G. Merlati, F. Valentino, C. Benatti, P. Gobbi, E. Ascari, "Avascular jaw osteonecrosis associated with bisphosphonate therapy," *Recenti Prog Med.*, vol. 97, no. 3, pp. 140-144, 2006.



- [39] P. Vescovi, E. Merigo, M. Meleti, M. Manfredi, "Bisphosphonate-associated osteonecrosis (BON) of the jaws: a possible treatment?," *J Oral Maxillofac Surg.*, vol. 64, no. 9, pp. 1460-1462, 2006.
- [40] L. Cavanna, R. Bertè, A. Arcari, P. Mordenti, R. Pagani, D. Vallisa, "Osteonecrosis of the jaw. A newly emerging site-specific osseous pathology in patients with cancer treated with bisphosphonates. Report of five cases and review of the literature," *Eur J Intern Med.*, vol. 18, no. 5, pp. 417-422, 2007.
- [41] R. Longo, M.A. Castellana, G. Gasparini, "Bisphosphonate-related osteonecrosis of the jaw and left thumb," *J Clin Oncol.*, vol. 27, no. 35, pp. 242-243, 2009.
- [42] A. Del Conte, P. Bernardeschi, F. La Ferla, G. Turrisi, M. D'Alessandro, F. Montagnani, G. Fiorentini, "Bisphosphonate-induced osteonecrosis of the jaw 32 months after interruption of zoledronate in a patient with multiple myeloma," *J Oral Maxillofac Surg.*, vol. 68, no. 5, pp. 1179-1782, 2010.
- [43] S. Papagni, N. Mongelli, M. Cozzolino, D. Brancaccio, F. Ciccolella, G. Favia, "Avascular jaw osteonecrosis in a hemodialysis patient treated with bisphosphonates," *G Ital Nefrol.*, vol. 24, no. 3, pp. 230-234, 2007.
- [44] E. Nastro, A. Allegra, G. Oteri, S. Ferlito, A. Alonci, G. Bellomo, A. D'Angelo, A. Tolomeo, D. Ciccì, F.S. De Ponte, C. Musolino, "Avascular necrosis of bone in leukemia and osteonecrosis of jaw by bisphosphonates," *J Oral Maxillofac Surg.*, vol. 67, no. 12, pp. 2701-2703, 2009.
- [45] A. Sardella, A. Carrassi, M. Tarozzi, G. Lodi, "Bisphosphonate-related osteonecrosis of the jaws associated with photodynamic therapy," *J Oral Maxillofac Surg.*, vol. 69, no. 10, pp. 314-316, 2011.
- [46] V. Fusco, A. Loidoris, G. Colella, P. Vescovi, G. Campisi, "Osteonecrosis of the jaw (ONJ) risk in breast cancer patients after zoledronic acid treatment," *Breast*, vol. 19, no. 5, pp. 432-433, 2010.
- [47] V. Guarneri, D. Miles, N. Robert, V. Diéras, J. Glaspy, I. Smith, C. Thomssen, L. Biganzoli, T. Taran, P. Conte, "Bevacizumab and osteonecrosis of the jaw: incidence and association with bisphosphonate therapy in three large prospective trials in advanced breast cancer," *Breast Cancer Res Treat.*, vol. 122, no. 1, pp. 181-188, 2010.
- [48] C. Ortega, F. Montemurro, R. Faggiuolo, R. Vormola, D. Nanni, F. Goia, M.O. Gilardino, M. Aglietta, "Osteonecrosis of the jaw in prostate cancer patients with bone



metastases treated with zoledronate: a retrospective analysis," *Acta Oncol.*, vol. 46, no. 5, pp. 664-668, 2007.

[49] A. Berruti, C. Ortega, V. Fusco, "Re: Christan Walter, Bilal Al-Nawas, Knut A. Grötz, et al. Prevalence and risk factors of bisphosphonate-associated osteonecrosis of the jaw in prostate cancer patients with advanced disease treated with zoledronate," *Eur Urol.*, vol. 55, no. 3, pp. 61-62, 2009.

[50] P. Tosi, E. Zamagni, D. Cangini, P. Tacchetti, F. Di Raimondo, L. Catalano, A. D'Arco, S. Ronconi, C. Cellini, M. Offidani, G. Perrone, M. Ceccolini, A. Brioli, S. Tura, M. Baccarani, M. Cavo, "Osteonecrosis of the jaws in newly diagnosed multiple myeloma patients treated with zoledronic acid and thalidomide-dexamethasone," *Blood.*, vol. 108, no. 12, pp. 3951-3952, 2006.

[51] S. Pozzi, R. Marcheselli, S. Sacchi, L. Baldini, F. Angrilli, E. Pennese, G. Quarta, C. Stelitano, G. Caparotti, S. Luminari, P. Musto, D. Natale, C. Broglia, A. Cuoghi, D. Dini, P. Di Tonno, G. Leonardi, G. Pianezze, V. Pitini, G. Polimeno, L. Ponchio, L. Masini, M. Musso, M. Spriano, G. Pollastri, "Bisphosphonate-associated osteonecrosis of the jaw: a review of 35 cases and an evaluation of its frequency in multiple myeloma patients," *Leuk Lymphoma.*, vol. 48, no. 1, pp. 56-64, 2007.

[52] E. Nastro, C. Musolino, A. Allegra, G. Oteri, M. Cicciù, A. Alonci, E. Quartarone, C. Alati, F.S. De Ponte, "Bisphosphonate-associated osteonecrosis of the jaw in patients with multiple myeloma and breast cancer," *Acta Haematol.* vol. 117, no. 3, pp. 181-187, 2006.

[53] A.M. Cafro, L. Barbarano, A.M. Nosari, G. D'Avanzo, M. Nichelatti, M. Bibas, D. Gaglioti, A. Taroni, F. Riva, E. Morra, A. Andriani, "Osteonecrosis of the jaw in patients with multiple myeloma treated with bisphosphonates: definition and management of the risk related to zoledronic acid," *Clin Lymphoma Myeloma*, vol. 8, no. 4, pp. 260, 2008.

[54] S. Pozzi, R. Marcheselli, S. Falorio, L. Masini, C. Stelitano, A. Falcone, G. Quarta, L. Ponchio, V. Pitini, S. Luminari, L. Baldini, "Bisphosphonates-associated osteonecrosis of the jaw: A long-term follow-up of a series of 35 cases observed by GISL and evaluation of its frequency over time," *Am J Hematol.* , vol. 84, no. 12, pp. 850-852, 2009.

[55] N. La Verde, C. Bareggi, M. Garassino, K. Borgonovo, P. Sburlati, D. Pedretti, C. Bianchi, S. Perrone, D. Mihali, S. Cobelli, C. Mantica, A. Rizzo, G. Farina, "Osteonecrosis of the jaw (ONJ) in cancer patients treated with Bisphosphonates: how the knowledge of a phenomenon can change its evolution," *Support Care Cancer.*, vol. 16, no. 11, pp. 1311-1315, 2008.



- [56] V. Fusco, A. Baraldi, A. Loidoris, L. Ciuffreda, C. Galassi, F. Goia, M. Scoletta, S. Berrone, A. Benech, C. Ortega, A. Berruti, "Jaw osteonecrosis associated with intravenous bisphosphonate: is incidence reduced after adoption of dental preventive measures?," *J Oral Maxillofac Surg.*, vol. 67, no. 8, pp. 1775, 2009.
- [57] M. Bonomi, R. Nortilli, A. Molino, T. Sava, A. Santo, A. Caldara, G.L. Cetto, "Renal toxicity and osteonecrosis of the jaw in cancer patients treated with bisphosphonates: a long-term retrospective analysis," *Med Oncol.*, vol. 27, no. 2, pp. 224-229, 2009.
- [58] P. Vescovi, E. Merigo, M. Meleti, M. Manfredi, R. Guidotti, S. Nammour, "Bisphosphonates-related osteonecrosis of the jaws: a concise review of the literature and a report of a single-centre experience with 151 patients," *J Oral Pathol Med.*, 2011.
- [59] S. Ferlito, S. Puzzo, F. Palermo, P. Verzi, "Treatment of bisphosphonate-related osteonecrosis of the jaws: presentation of a protocol and an observational longitudinal study of an Italian series of cases," *Br J Oral Maxillofac Surg.*, 2011 Sep.
- [60] V. Fusco, C. Ortega, G. Gorzegno, A. Loidoris, A. Baraldi, A. Vandone, A. Fasciolo, E. Pomatto, M. Migliario, O. Bertetto. "Increase and decrease of jaw osteonecrosis (ONJ) in patients treated with intravenous bisphosphonates (BP): impact of preventive measures and reduced prescriptions in the experience of the "Rete Oncologica di Piemonte e Valle d'Aosta" ONJ study group. *European Journal of Cancer Supplements*, Vol 7 No 2, September 2009, Page 201
- [61] P. Vescovi, G. Campisi, V. Fusco, G. Mergoni, M. Manfredi, E. Merigo, L. Solazzo, M. Gabriele, G.M. Gaeta, G. Favia, F. Peluso, G. Colella, "Surgery-triggered and non surgery-triggered Bisphosphonate-related Osteonecrosis of the Jaws (BRONJ): A retrospective analysis of 567 cases in an Italian multicenter study," *Oral Oncol.*, vol. 47, no. 3, pp. 191-194, 2011.
- [62] G. Colella, G. Campisi, V. Fusco, "American Association of Oral and Maxillofacial Surgeons position paper: Bisphosphonate-Related Osteonecrosis of the Jaws-2009 update: the need to refine the BRONJ definition," *J Oral Maxillofac Surg.*, vol. 67, no. 12, pp. 2698-2699, 2009.
- [63] S. Fedele, S.R. Porter, F. D'Aiuto, S. Aljohani, P. Vescovi, M. Manfredi, P.G. Arduino, R. Broccoletti, A. Musciotto, O. Di Fede, T.S. Lazarovici, G. Campisi, N. Yarom, "Nonexposed variant of bisphosphonate-associated osteonecrosis of the jaw: a case series," *Am J Med.*, vol. 123, no. 11, pp. 1060-1064, 2010



- [64] S. Chiandussi, M. Biasotto, F. Dore, F. Cavalli, M.A. Cova, R. Di Lenarda, "Clinical and diagnostic imaging of bisphosphonate-associated osteonecrosis of the jaws," *Dentomaxillofac Radiol.*, vol. 35, no. 4, pp. 236-243, 2006.
- [65] S.D. Bianchi, M. Scoletta, F.B. Cassione, G. Migliaretti, M. Mozzati, "Computerized tomographic findings in bisphosphonate-associated osteonecrosis of the jaw in patients with cancer," *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.*, vol. 104, no. 2, pp. 249-258, 2007.
- [66] L. Catalano, S. Del Vecchio, F. Petruzzello, R. Fonti, B. Salvatore, C. Martorelli, C. Califano, G. Caparrotti, S. Segreto, L. Pace, "Sestamibi and FDG-PET scans to support diagnosis of jaw osteonecrosis," *Ann Hematol.* vol. 86, no. 6, pp. 415-423, 2007.
- [67] P. Milillo, A.P. Garribba, G. Favia, G.C. Ettore, "Jaw osteonecrosis in patients treated with bisphosphonates: MDCT evaluation," *Radiol Med.*, vol. 112, no. 4, pp. 603-611, 2007.
- [68] A. Bedogni, S. Blandamura, Z. Lokmic, C. Palumbo, M. Ragazzo, F. Ferrari, A. Tregnaghi, F. Pietrogrande, O. Procopio, G. Saia, M. Ferretti, G. Bedogni, L. Chiarini, G. Ferronato, V. Ninfo, L. Lo Russo, L. Lo Muzio, P.F. Nocini, "Bisphosphonate-associated jawbone osteonecrosis: a correlation between imaging techniques and histopathology," *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.*, vol. 105, no. 3, pp. 358-364, 2008.
- [69] F. Dore, L. Filippi, "Bone Scintigraphy and SPECT/CT in Bisphosphonate-Induced Osteonecrosis of the Jaw," *J Nucl Med.*, 2009.
- [70] D. Carmagnola, S. Celestino, S. Abati, "Dental and periodontal history of oncologic patients on parenteral bisphosphonates with or without osteonecrosis of the jaws: a pilot study," *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.*, vol. 106, no. 6, pp. 10-15, 2008.
- [71] R. Addeo, V. Nocera, V. Faiola, B. Vincenzi, G. Ferraro, L. Montella, R. Guarrasi, E. Rossi, G. Cennamo, G. Tonini, E. Capasso, D. Santini, M. Caraglia, S. Del Prete, "Management of pain in elderly patients receiving infusion of zoledronic acid for bone metastasis: a single-institution report," *Support Care Cancer.*, vol. 16, no. 2, pp. 209-214, 2008.
- [72] A. Corso, M. Varettoni, P. Zappasodi, C. Klersy, S. Mangiacavalli, G. Pica, M. Lazzarino, "A different schedule of zoledronic acid can reduce the risk of the osteonecrosis of the jaw in patients with multiple myeloma," *Leukemia.*, vol. 27, no. 7, pp. 1545-1548, 2007.



- [73] A. Bedogni, G. Bettini, A. Totola, G. Saia, P.F. Nocini, "Oral bisphosphonate-associated osteonecrosis of the jaw after implant surgery: a case report and literature review," *J Oral Maxillofac Surg.*, vol. 68, no. 7, pp. 1662-1666, 2010.
- [74] G. Favia, A. Piattelli, P. Sportelli, S. Capodiferro, G. Iezzi, "Osteonecrosis of the posterior mandible after implant insertion: a clinical and histological case report," *Clin Implant Dent Relat Res.*, vol. 13, no. 1, pp. 58-63, 2011.
- [75] S. Ferlito, C. Liardo, S. Puzzo, "Dental extractions in patient treated with intravenous bisphosphonates and risk of osteonecrosis of jaws: presentation of a preventive protocol and case series," *Minerva Stomatol.*, vol. 59, no. 11-12, pp. 593-601, 2010.
- [76] M. Biasotto, S. Chiandussi, F. Dore, A. Rinaldi, C. Rizzardi, F. Cavalli, R. Di Lenarda, "Clinical aspects and management of bisphosphonates-associated osteonecrosis of the jaws," *Acta Odontol Scand.*, vol. 64, no. 3, pp. 348-354, 2006.
- [77] F. Capsoni, M. Longhi, R. Weinstein, "Bisphosphonate-associated osteonecrosis of the jaw: the rheumatologist's role," *Arthritis Res Ther.*, vol. 8, no. 5, pp. 219, 2006.
- [78] G. Ficarra, F. Beninati, "Bisphosphonate-related osteonecrosis of the jaws: an update on clinical, pathological and management aspects," *Head Neck Pathol.*, vol. 1, no. 2, pp. 132-140, 2007.
- [79] G. Campisi, O. Di Fede, A. Musciotto, A. Lo Casto, L. Lo Muzio, F. Fulfaro, G. Badalamenti, A. Russo, N. Gebbia, "Bisphosphonate-related osteonecrosis of the jaw (BRONJ): current dental management designs and issues in diagnosis," *Ann Oncol.*, vol. 18, Suppl. 6, pp. 68-72, 2007.
- [80] F. Capsoni, "[Bisphosphonate-associated osteonecrosis of the jaw in rheumatology: a systematic review]," *Reumatismo*, vol. 60, no. 1, pp. 6-13, 2008.
- [81] V. Montefusco, F. Gay, F. Spina, R. Miceli, M. Maniezzo, M. Teresa Ambrosini, L. Farina, S. Piva, A. Palumbo, M. Boccadoro, P. Corradini, "Antibiotic prophylaxis before dental procedures may reduce the incidence of osteonecrosis of the jaw in patients with multiple myeloma treated with bisphosphonates," *Leuk Lymphoma.*, Vol. 49, no. 11, pp. 2156-2162, 2008.
- [82] G. Campisi, S. Fedele, G. Colella, A.L. Casto, V. Fusco, "Canadian consensus practice guidelines for bisphosphonate associated osteonecrosis of the jaw," *J Rheumatol.*, vol. 36, no. 2, pp. 451-453, 2009.



- [83] S. Fedele, N. Kumar, R. Davies, J. Fiske, S. Greening, S. Porter, "Dental management of patients at risk of osteonecrosis of the jaws: a critical review," *Oral Dis.*, vol. 15, no. 8, pp. 527-537, 2009.
- [84] A. Borgioli, C. Viviani, M. Duvina, L. Brancato, G. Spinelli, M.L. Brandi, P. Tonelli, "Biphosphonates-related osteonecrosis of the jaw: Clinical and physiopathological considerations," *Clin Risk Manag.*, vol. 5, no. 1, pp. 217-227, 2009.
- [85] C.I. Ripamonti, M. Maniezzo, T. Campa, E. Fagnoni, C. Brunelli, G. Saibene, C. Bareggi, L. Ascani, E. Cislighi, "Decreased occurrence of osteonecrosis of the jaw after implementation of dental preventive measures in solid tumour patients with bone metastases treated with bisphosphonates. The experience of the National Cancer Institute of Milan," *Ann Oncol.*, vol. 20, no. 1, pp. 137-145, 2009.
- [86] A.M. Vandone, M. Donadio, M. Mozzati, M. Ardine, M.A. Polimeni, S. Beatrice, L. Ciuffreda, M. Scoletta, "Impact of dental care in the prevention of bisphosphonate-associated osteonecrosis of the jaw: a single-center clinical experience," *Ann Oncol.*, 2011.
- [87] G. Lodi, A. Sardella, A. Salis, F. Demarosi, M. Tarozzi, A. Carrassi, "Tooth extraction in patients taking intravenous bisphosphonates: a preventive protocol and case series," *J Oral Maxillofac Surg.*, vol. 68, no. 1, pp. 107-110, 2010.
- [88] M. Scoletta, P.G. Arduino, R. Pol, V. Arata, S. Silvestri, A. Chiecchio, M. Mozzati, "Initial experience on the outcome of teeth extractions in intravenous bisphosphonate-treated patients: a cautionary report," *J Oral Maxillofac Surg.*, vol. 69, no. 2, pp. 456-462, 2010.
- [89] S. Ferlito, S. Puzzo, C. Liardo, "Preventive protocol for tooth extractions in patients treated with zoledronate: a case series," *J Oral Maxillofac Surg.*, vol. 69, no. 6, pp. 1-4, 2011.
- [90] F. Bertoldo, G. Bianchi, "[Comment on the article "Bisphosphonate-associated osteonecrosis of the jaw in rheumatology: a systematic review" by Capsoni Franco]," *Reumatismo*, vol. 60, no. 3, pp. 234-235, 2008.
- [91] F. Bertoldo, F. Aging, "Bisphosphonates and osteonecrosis of the mandible/maxilla in osteoporosis: no reason to panic," *Clin Exp Res.*, vol.20, no.1, pp. 87-90, 2008.
- [92] G. Favia, G.P. Pilolli, E. Maiorano, "Histologic and histomorphometric features of bisphosphonate-related osteonecrosis of the jaws: an analysis of 31 cases with confocal laser scanning microscopy," *Bone*, vol. 45, no. 3, pp. 406-413, 2009.
- [93] G. Gasparini, G. Saponaro, F. Di Nardo, A. Moro, R. Boniello, D. Cervelli, T.M. Marianetti, G. Palazzoni, S. Pelo, "Clinical experience with spiramycin in bisphosphonate-



- associated osteonecrosis of the jaw," *Int J Immunopathol Pharmacol.*, vol. 23, no. 2, pp. 619-626, 2010.
- [94] M. Manfredi, E. Merigo, R. Guidotti, M. Meleti, P. Vescovi, "Bisphosphonate-related osteonecrosis of the jaws: a case series of 25 patients affected by osteoporosis," *Int J Oral Maxillofac Surg.*, vol. 40, no. 3, pp. 277-284, 2011.
- [95] S. Otto, M.H. Abu-Id, S. Fedele, P.H. Warnke, S.T. Becker, A. Kolk, T. Mücke, G. Mast, R. Köhnke, E. Volkmer, F. Haasters, O. Lieger, T. Iizuka, S. Porter, G. Campisi, G. Colella, O. Ploder, A. Neff, J. Wiltfang, M. Ehrenfeld, T. Kreusch, K.D. Wolff, S.R. Stürzenbaum, M. Schieker, C. Pautke, "Osteoporosis and bisphosphonates-related osteonecrosis of the jaw: not just a sporadic coincidence--a multi-centre study," *J Craniomaxillofac Surg.*, vol. 39, no. 4, pp. 272-277, 2010.
- [96] E. Serra, M. Paolantonio, G. Spoto, F. Mastrangelo, S. Tetè, M. Dolci "Bevacizumab-related osteonecrosis of the jaw," *Int J Immunopathol Pharmacol.*, vol. 22, no. 4, pp. 1121-1123, 2009.
- [97] A. Brunello, G. Saia, A. Bedogni, D. Scaglione, U. Basso, "Worsening of osteonecrosis of the jaw during treatment with sunitinib in a patient with metastatic renal cell carcinoma," *Bone.*, vol. 44, no. 1, pp. 173-175, 2008.
- [98] V. Fusco, C. Porta, C. Paglino, A. Bedogni, G. Bettini, G. Saia, G. Campisi, M. Scoletta, A. Agrillo, A. Fasciolo "Osteonecrosis of the jaw (ONJ) in patients with renal cancer treated with bisphosphonates (BPs) and sunitinib or other targeted therapy (TT) agents: A multicenter survey" . *J Clin Oncol 29 (suppl):* 2011 (suppl; abstr e15182)
- [99] R. Diego, O. D'Orto, D. Pagani, A. Agazzi, U. Marzano, G. Derada Troletti, W. Fontanella, L. Pignataro, "Bisphosphonate-associated osteonecrosis of the jaws: a therapeutic dilemma," *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.*, vol. 103, no. 3, pp. 1-5, 2007.
- [100] M. Scoletta, P.G. Arduino, P. Dalmaso, R. Broccoletti, M. Mozzati, "Treatment outcomes in patients with bisphosphonate-related osteonecrosis of the jaws: a prospective study," *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.*, vol. 110, no. 1, pp. 46-53, 2010.
- [101] P. Vescovi, S. Nammour, "Bisphosphonate-Related Osteonecrosis of the Jaw (BRONJ) therapy. A critical review," *Minerva Stomatol.*, vol. 59, no. 4, pp. 181-203, 2010.



- [102] L. Montebugnoli, L. Felicetti, D.B. Gissi, A. Pizzigallo, G.A. Pelliccioni, C. Marchetti, "Biphosphonate-associated osteonecrosis can be controlled by nonsurgical management," *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.*, vol. 104, no. 4, pp. 473-477, 2007.
- [103] F. Moretti, G.A. Pelliccioni, L. Montebugnoli, C. Marchetti, "A prospective clinical trial for assessing the efficacy of a minimally invasive protocol in patients with bisphosphonate-associated osteonecrosis of the jaws," *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.*, vol. 112, no. 6, pp. 777-782, 2011.
- [104] G. Longobardi, R. Boniello, G. Gasparini, I. Pagano, S. Pelo, "Surgical therapy for osteonecrotic lesions of the jaws in patients in therapy with bisphosphonates," *J Craniofac Surg.*, vol. 18, no. 5, pp. 1012-1017, 2007.
- [105] P. Vescovi, M. Manfredi, E. Merigo, M. Meleti, "Early surgical approach preferable to medical therapy for bisphosphonate-related osteonecrosis of the jaws," *J Oral Maxillofac Surg.*, vol. 66, no. 4, pp. 831-832, 2008.
- [106] B. Bianchi, A. Ferri, S. Ferrari, C. Copelli, E. Sesenna, "Subtotal mandibular reconstruction using an intraoral approach: report of 2 cases," *J Oral Maxillofac Surg.*, vol. 66 no.12 pp.2654-2656, 2008.
- [107] S. Ferrari, B. Bianchi, A. Savi, T. Poli, A. Multinu, A. Balestreri, A. Ferri, "Fibula free flap with endosseous implants for reconstructing a resected mandible in bisphosphonate osteonecrosis," *J Oral Maxillofac Surg.*, vol. 66, no. 5, pp. 999-1003, 2008.
- [108] P.F. Nocini, G. Saia, G. Bettini, M. Ragazzo, S. Blandamura, L. Chiarini, A. Bedogni, "Vascularized fibula flap reconstruction of the mandible in bisphosphonate-related osteonecrosis," *Eur J Surg Oncol.*, vol. 35, no. 4, pp. 373-379, 2008.
- [109] B. Bianchi, C. Copelli, S. Ferrari, A. Ferri, E. Sasenna, "Free flaps: outcomes and complications in head and neck reconstructions," *J Craniomaxillofac Surg.*, vol.37 no.8, pp.438-42, 2009.
- [110] G. Tirelli, M. Biasotto, S. Chiandussi, F. Dore, E. De Nardi, R. Di Lenarda, "Bisphosphonate-associated osteonecrosis of the jaws: the limits of a conservative approach," *Head Neck.*, vol. 31, no. 9, pp. 1249-1254, 2009.
- [111] R. Sacco, G. Sacco, A. Acocella, S. Sale, N. Sacco, E. Baldoni, "A systematic review of microsurgical reconstruction of the jaws using vascularized fibula flap technique in patients with bisphosphonate-related osteonecrosis," *J Appl Oral Sci.*, vol. 19, no. 4, pp. 293-300, 2011.

- [112] A. Bedogni, G. Saia, G. Bettini, A. Tronchet, A. Totola, G. Bedogni, G. Ferronato, P.F. Nocini, S. Blandamura, "Long-term outcomes of surgical resection of the jaws in cancer patients with bisphosphonate-related osteonecrosis," *Oral Oncol.*, vol. 47, no. 5, pp. 420-440, 2011.
- [113] L. Cella, A. Oppici, M. Arbasi, M. Moretto, M. Piepoli, D. Vallisa, A. Zangrandi, C. Di Nunzio, L. Capanna, "Autologous bone marrow stem cell intralesional transplantation repairing bisphosphonate related osteonecrosis of the jaw," *Head Face Med.*, vol. 17, no. 7, pp. 16, 2011.
- [114] P. Vescovi, E. Merigo, M. Meleti, C. Fornaini, S. Nammour, M. Manfredi, "Nd:YAG laser biostimulation of bisphosphonate-associated necrosis of the jawbone with and without surgical treatment," *Br J Oral Maxillofac Surg.*, vol. 45, no. 8, pp. 628-632, 2007.
- [115] P. Vescovi, E. Merigo, M. Manfredi, M. Meleti, C. Fornaini, M. Bonanini, J.P. Rocca, S. Nammour, "Nd:YAG laser biostimulation in the treatment of bisphosphonate-associated osteonecrosis of the jaw: clinical experience in 28 cases," *Photomed Laser Surg.*, vol. 26, no. 1, pp. 37-46, 2008.
- [116] F. Angiero, C. Sannino, R. Borloni, R. Crippa, S. Benedicenti, G.E. Romanos, "Osteonecrosis of the jaws caused by bisphosphonates: evaluation of a new therapeutic approach using the Er:YAG laser," *Lasers Med Sci.*, vol. 24, no. 6, pp. 849-856, 2009.
- [117] P. Vescovi, M. Manfredi, E. Merigo, M. Meleti, C. Fornaini, J.P. Rocca, S. Nammour "Surgical approach with Er:YAG laser on osteonecrosis of the jaws (ONJ) in patients under bisphosphonate therapy (BPT)," *Lasers Med Sci.*, vol. 25, no. 1, pp. 101-103, 2009.
- [118] M. Scoletta, P.G. Arduino, L. Reggio, P. Dalmasso, M. Mozzati, "Effect of low-level laser irradiation on bisphosphonate-induced osteonecrosis of the jaws: preliminary results of a prospective study," *Photomed Laser Surg.*, vol. 28, no. 2, pp. 179-184, 2010.
- [119] U. Romeo, A. Galanakis, C. Marias, A.D. Vecchio, G. Tenore, G. Palaia, P. Vescovi, A. Polimeri, "Observation of pain control in patients with bisphosphonate-induced osteonecrosis using low level laser therapy: preliminary results," *Photomed Laser Surg.*, vol. 29, no. 7, pp. 447-452, 2011.
- [120] A. Agrillo, M.T. Petrucci, M. Tedaldi, M.C. Mustazza, S.M. Marino, C. Gallucci, G. Iannetti, "New therapeutic protocol in the treatment of avascular necrosis of the jaws," *J Craniofac Surg.*, vol. 17, no. 6, pp. 1080-1083, 2006.



- [121] M.T. Petrucci, C. Gallucci, A. Agrillo, M.C. Mustazza, R. Foà, "Role of ozone therapy in the treatment of osteonecrosis of the jaws in multiple myeloma patients," *Haematologic*, vol. 92, no. 9, pp. 1289-1290, 2007.
- [122] A. Agrillo, C. Ungari, F. Filiaci, P. Priore, G. Iannetti, "Ozone therapy in the treatment of avascular bisphosphonate-related jaw osteonecrosis," *J Craniofac Surg.*, vol. 18, no. 5, pp. 1071-1075, 2007
- [123] C.I. Ripamonti, E. Cislighi, L. Mariani, M. Maniezzo, "Efficacy and safety of medical ozone (O₃) delivered in oil suspension applications for the treatment of osteonecrosis of the jaw in patients with bone metastases treated with bisphosphonates: Preliminary results of a phase I-II study," *Oral Oncol.*, vol 47, no. 3, pp. 185-190, 2011.
- [124] P.Vescovi, "Osteonecrosi mascellari e bifosfonati: prevenzione, diagnosi e terapia," *Stamperia*, Parma, Italia, 2007.
- [125] P.Vescovi, "Osteonecrosi dei mascellari e bifosfonati" *Tecniche Nuove*, Milano, Italia, 2008.
- [126] R.E.Marx, "L' osteonecrosi dei mascellari da bifosfonati - Storia, Eziologia, Prevenzione e trattamento dell' osteonecrosi indotta dai bifosfonati per via orale e per via endovenosa" *Quintaessenza*, Rho, Italia, 2009 (translated by S.Abati et al) .
- [127] L.Sottosanti " Le segnalazioni all'AIFA. I dati della Rete Nazionale di Farmacovigilanza (RNF) sui Bifosfonati e ONJ " in "Osteonecrosi dei mascellari (ONJ): Prevenzione, Diagnosi, Trattamento. Update 2009" available at http://www.reteoncologica.it/images/stories/ONJ/Osteonecrosi_mascellari_2009/02-04%20Sottosanti%20Laura.pdf
- [128] EMEA "CHMP assessment report on bisphosphonates and osteonecrosis of the jaw". London, 24 September 2009, available at http://www.emea.europa.eu/docs/en_GB/document_library/Report/2010/01/WC500051428.pdf
- [129] L.Sottosanti " Osteonecrosi della mandibola e farmaci biologici " in "Reazioni. Bollettino di farmacovigilanza dell'AIFA" vol 14 (settembre 2009) pp 6-7 , available at http://www.sefap.it/farmacovigilanza_news_200912/rea_14_2009.pdf
- [130] Additional safety information. In *European Medicines Agency Monthly report* (26 November 2010), pag 2, available at http://www.ema.europa.eu/docs/en_GB/document_library/Committee_meeting_report/2010/11/WC500099368.pdf



- [131] AIFA. "Nota Informativa Importante su Avastin (bevacizumab) del 30/11/2010" available at http://www.agenziafarmaco.gov.it/sites/default/files/nota_informativa_importante_sullavastin_del_30-11-2010.pdf
- [132] AIFA. "Nota Informativa Importante su Sunitinib (Sutent) del 30/11/2010" available at http://www.agenziafarmaco.gov.it/sites/default/files/nota_informativa_importante_su_sunitinib_30-11-10.pdf
- [133] Gruppo di Lavoro sull'Osteonecrosi da Bifosfonati della Rete Oncologica di Piemonte e Valle d'Aosta "Newsletter n5 (Dicembre 2009)", available at http://www.reteoncologica.it/images/stories/scaffale/archivio/Gruppo_Osteonecrosi_Newsletter_5_2009.pdf
- [134] Osteonecrosi dei mascellari : "Slide gallery", available at http://www.reteoncologica.it/index.php?option=com_content&view=article&id=134&Itemid=85
- [135] Ministero del Lavoro, della Salute e delle Politiche Sociali "Raccomandazione per la prevenzione dell'osteonecrosi della mascella / mandibola da bifosfonati" Raccomandazione n.10, settembre 2009, available at http://www.salute.gov.it/imgs/C_17_pubblicazioni_1077_allegato.pdf
- [136] V. Fusco, C. Ortega, F. Goia, L. Ciuffreda, M. Ardine, P. Falco, G. Gorzeglio, A. Baraldi, M. Airoidi, O. Bertetto "Bisphosphonates (BP)-related Osteonecrosis of the Jaw (ONJ): clinical experience of the "Rete Oncologica di Piemonte e Valle d'Aosta" network." *Journal of Clinical Oncology*, 2006 ASCO Annual Meeting Proceedings Part I. Vol 24, No. 18S (June 20 Supplement): 18584, 2006
- [137] N. D'Alessandro, I. Morreale "Esperienza della Regione Sicilia" in in "Osteonecrosi dei mascellari (ONJ): Prevenzione, Diagnosi, Trattamento. Update 2009", available at <http://www.reteoncologica.it/images/stories/slideOnline/osteonecrosi/morreale%20-%20dalessandro.pdf>
- [138] BE Hillner, JN Ingle, JR Berenson, et al: American Society of Clinical Oncology Guideline on the Role of Bisphosphonates in Breast Cancer. *J Clin Oncol* 18:1378-1391, 2000
- [139] BE Hillner, JN Ingle, RT Chlebowski, et al: American Society of Clinical Oncology 2003 update on the role of bisphosphonates and bone health issues in women with breast cancer. *J Clin Oncol* 21:4042-4057, 2003



- [140] JR Berenson et al "American Society of Clinical Oncology clinical practice guidelines: the role of bisphosphonates in multiple myeloma". *J Clin Oncol* 20(17): pp3719-36, 2002
- [141] M.Venegoni "Ritorna, coi bifosfonati nuovi, la vecchia necrosi dei fiammiferai" *Occhio clinico* n. 6, pp 15-16, 2005
- [142] BG Durie, M Katz, J Crowley."Osteonecrosis of the jaw and bisphosphonates" *N Engl J Med.* 353(1): pp 99-102, 2005 .
- [143] M Zambelli, M Scoletta, M Mozzati "L'informazione del paziente come elemento di prevenzione della BRONJ" in "*Osteonecrosi dei mascellari (ONJ): Prevenzione, Diagnosi, Trattamento. UPDATE 2010*", pp 43, 2010 , available at http://www.reteoncologica.it/index.php?option=com_content&view=article&id=134&Itemid=85
4. [144] CA Migliorati, K Mattos, MJ Palazzolo. "How patients' lack of knowledge about oral bisphosphonates can interfere with medical and dental care". *J Am Dent Assoc.* 141(5):562-6, 2010
5. [145] JS Bauer, N Beck, J Kiefer, P Stockmann, M Wichmann, S Eitner. "Awareness and education of patients receiving bisphosphonates". *J Craniomaxillofac Surg.* , 2011 (epub)
6. [146] M Aapro, PA Abrahamsson, JJ Body, RE Coleman, R Colomer, L Costa, L Crinò, L Dirix, M Gnant, J Gralow, P Hadji, GN Hortobagyi, W Jonat, A Lipton, A Monnier, AH Paterson, R Rizzoli , F Saad, B Thürlimann. "Guidance on the use of bisphosphonates in solid tumours: recommendations of an international expert panel" *Ann Oncol.* 19(3): pp 420-32, 2008
7. [147] RA Kyle, GC Yee, MR Somerfield, PJ Flynn, S Halabi, S Jagannath, RZ Orlowski, DG Roodman, P Twilde, K Anderson; American Society of Clinical Oncology. "American Society of Clinical Oncology 2007 clinical practice guideline update on the role of bisphosphonates in multiple myeloma". *J Clin Oncol.* 25(17): pp2464-72, 2007

FIGURES




Figure 1: Number of prescribed units of pamidronate (1 unit = 90 mg) and zoledronic acid (1 unit = 4 mg) in Italy between 2001 and 2008 (elaboration from AIFA data, personal communication).




Figure 2: Rate of prescribed units of zoledronic acid (1 unit = 4 mg) in Italy between 2001 and 2006, per million of inhabitants, in several European countries (elaboration from AIFA data, personal communication).